

Tabulation of PVI Transcendents and Parametrization Formulas

Davide Guzzetti¹

First submitted to arXiv on August 17, 2011

arXiv 1108.3401v1

Abstract

The critical and asymptotic behaviors of solutions of the sixth Painlevé equation PVI, obtained in the framework of the monodromy preserving deformation method, and their explicit parametrization in terms of monodromy data, are synthetically reviewed and tabulated.

MSC: 34M55 (Painlevé and other special functions)

1 Introduction

In the last few decades, the Painlevé equations have emerged as one of the central objects in pure mathematics and mathematical physics. They define new non-linear special functions, which find applications in a variety of problems, such as number theory, theory of analytic varieties (like Frobenius structures), random matrix theory, orthogonal polynomials, non linear evolutionary PDEs, combinatorial problems, etc. The properties of the classical special functions have been organized and tabulated in various classical handbooks. The scientific community is now engaged in the project of a comparable organization and tabulation of the properties of the Painlevé functions (this is part of the *Painlevé Project*, see <http://cio.nist.gov/esd/emaildir/lists/painleveproject/>).

In this paper, I provide two tables, one for the critical behaviors of solutions of the sixth Painlevé equation PVI, and one for the formulas which parameterize, in terms of associated monodromy data, the integration constants determining the critical behaviors.

Though critical behaviors and expansions of the Painlevé six functions at the critical points, and their parameterization in terms of monodromy data, have been extensively studied, the effort of collecting and tabulating them has been missing. It is the time to make a first point of the situation, in view of the Painlevé Project. This is the aim of the paper.

The tables are based on the results of Jimbo [17] and of my papers [10], [11], [12] and [9]. They represent essentially all have been obtained about the critical behaviors of the Painlevé six functions and their parameterization in terms of associated monodromy data, by techniques of the monodromy preserving deformations method.

The table of critical behaviors I propose here is essentially complete. Namely, the solutions which are tabulated form a set in correspondence to (essentially) the whole space of monodromy data of the isomonodromic Fuchsian system associated to PVI.

Except for the case when logarithms appear in the expansions tabulated, the expansions are convergent and define true solutions. This follows from the results of [22] and [9]. When logarithms appear, no proof of convergence is known to the author, though one may expect that an extension of the method of [22] and [9] be possible for the power logarithmic expansions.

¹) International School of Advanced Studies SISSA/ISAS, Via Bonomea 265, 34136 Trieste, Italy. E-mail: guzzetti@sissa.it

²) Korea Institute of Advanced Study KIAS, Hoegiro 87(207-43 Cheongnyangni-dong), Dongdaemun-gu, Seoul 130-722, Korea.

Bruno and Goryuchkina, in a series of papers which are summarized in their review [6], have computed the expansions of actual and possibly formal solutions of PVI. They have obtained all the expansions which can be constructed by means of their method of power geometry (see [4] and [5]). In section 7, I compare their results with those obtained with the method of monodromy preserving deformations, showing that they coincide.

The solutions (22) and (26) in the table of critical behaviors are also obtained by Shimomura [22], on the universal covering of a punctured neighborhood of the critical point.

The sixth Painlevé equation PVI is:

$$\begin{aligned} \frac{d^2 y}{dx^2} = & \frac{1}{2} \left[\frac{1}{y} + \frac{1}{y-1} + \frac{1}{y-x} \right] \left(\frac{dy}{dx} \right)^2 - \left[\frac{1}{x} + \frac{1}{x-1} + \frac{1}{y-x} \right] \frac{dy}{dx} \\ & + \frac{y(y-1)(y-x)}{x^2(x-1)^2} \left[\alpha + \beta \frac{x}{y^2} + \gamma \frac{x-1}{(y-1)^2} + \delta \frac{x(x-1)}{(y-x)^2} \right], \quad (\text{PVI}). \end{aligned}$$

Following [7] (page 8), *solving* PVI means that:

i) We determine the *explicit* critical behavior (or asymptotic expansion) of $y(x)$, by an explicit formula in terms of two integration constants.

ii) We solve the *connection problem*, namely we find the *explicit* relations among couples of integration constants at different critical points.

If we can solve i) and ii), then Painlevé transcendents can be efficiently used in applications as special functions. The tables of the critical behaviors and their parameterization formulas in terms of monodromy data give answer to i) and ii).²

In the following, the subject "I" will be replaced by "we". Then, we recall the basic fact that, according to [18], PVI is the isomonodromy deformation equation of the 2×2 Fuchsian system below:

$$\frac{d\Psi}{d\lambda} = A(x, \lambda) \Psi, \quad A(x, \lambda) := \left[\frac{A_0(x)}{\lambda} + \frac{A_x(x)}{\lambda-x} + \frac{A_1(x)}{\lambda-1} \right], \quad \lambda \in \mathbf{C}. \quad (1)$$

where $A(x, \lambda)$ depends on x in such a way that

$$A_{12}(x, \lambda) = \frac{f(x)(\lambda - y(x))}{\lambda(\lambda-1)(\lambda-x)}$$

Therefore,

$$y(x) = \frac{x(A_0)_{12}}{x[(A_0)_{12} + (A_1)_{12}] - (A_1)_{12}} \quad (2)$$

The 2×2 matrices $A_i(x)$ depend on x in such a way that there exists a fundamental matrix solution $\Psi(\lambda, x)$ with monodromy independent of small deformations of x . They also depend algebraically on $\alpha, \beta, \gamma, \delta$ according to the following relations:

$$A_0 + A_1 + A_x = -\frac{\theta_\infty}{2} \sigma_3, \quad \theta_\infty \neq 0. \quad \text{Eigenvalues } (A_i) = \pm \frac{1}{2} \theta_i, \quad i = 0, 1, x; \quad (3)$$

$$\alpha = \frac{1}{2}(\theta_\infty - 1)^2, \quad -\beta = \frac{1}{2}\theta_0^2, \quad \gamma = \frac{1}{2}\theta_1^2, \quad \left(\frac{1}{2} - \delta \right) = \frac{1}{2}\theta_x^2 \quad (4)$$

In the " λ -plane" $\mathbf{C} \setminus \{0, x, 1\}$ we fix a base point λ_0 and three loops, which are numbered in order 1, 2, 3 according to a counter-clockwise order referred to λ_0 . We choose $0, x, 1$ to be the order 1, 2, 3. We denote the loops by $\gamma_0, \gamma_x, \gamma_1$. See figure 1. The monodromy matrices of a fundamental solution

²A more restrictive definition of "solving" should include the determination of the distribution of the movable poles, for which we refer to [13] and [3].




Figure 1: The ordered basis of loops

$\Psi(\lambda)$ w.r.t. this base of loops are denoted M_0, M_x, M_1 . The loop at infinity will be $\gamma_\infty = \gamma_0\gamma_x\gamma_1$, so $M_\infty = M_1M_xM_0$. When λ goes around a small loop around $\lambda = i, i = 0, x, 1$, the fundamental solution transforms like $\Psi \mapsto \Psi M_i$. M_0, M_x, M_1 are the monodromy matrices. There is a *one-to-one correspondence between a point in the space of monodromy data and a branch $y(x)$* (except when one M_i or $M_1M_xM_0$ is the identity. See [11] and [12]). As a consequence, *the integration constants of $y(x)$ can be parameterized in terms of the monodromy data*. Jimbo provided for the first time the critical behavior and its parameterization in terms of monodromy data for a wide class of two complex parameter solutions. His work [17] is the foundation of all subsequent developments.

Organization of the paper

Section 2. The table of critical behaviors is provided.

Section 3. The table of the parametrization of basic critical behaviors in terms of monodromy data is provided.

Section 4. The completeness of the table of the critical behaviors is discussed.

Section 5. Okamoto's transformations are reviewed and the basic solutions are defined.

Section 6. The computation of critical behaviors through the matching procedure is reviewed.

Section 7. The table of critical behaviors is discussed in details. A comparison with the results of [6] is provided.

Section 8. Details on the parametrization of the critical behaviors in terms of monodromy data are given. The action of the Okamoto's transformations on the monodromy data is briefly reviewed.

Section 9. The computation of monodromy data through the matching procedure is reviewed.

2 Table of Critical Behaviors

Here the table is given of the critical behaviors of branches of solutions of PVI at $x = 0$. The branch cut may be taken to be $-\pi < \arg x \leq \pi$. This is the first tabulation available in the literature. The detailed description and explanation of the critical behaviors tabulated below is in section 7. The numeration of the expansions in the table is that of section 7. The term "basic solutions" is explained in section 5. In order to understand the table, we introduce the set $\Sigma_{\beta\delta} = \Sigma_{\beta\delta}^1 \cup \Sigma_{\beta\delta}^2$, where:

$$\Sigma_{\beta\delta}^k := \left\{ \left(\sqrt{-2\beta} + (-)^k \sqrt{1-2\delta} \right) \operatorname{sgn} \left(\Re \left(\sqrt{-2\beta} + \sqrt{1-2\delta} \right) \right) \right\}, \quad k = 1, 2, \quad (5)$$

or, in case $|\sqrt{-2\beta} + (-)^k \sqrt{1-2\delta}| < 1$,

$$\Sigma_{\beta\delta}^k := \left\{ \left(\sqrt{-2\beta} + (-)^k \sqrt{1-2\delta} \right), -\left(\sqrt{-2\beta} + (-)^k \sqrt{1-2\delta} \right) \right\}. \quad (6)$$

In the monodromy deformation parameters, this is $\Sigma_{\beta\delta} = \{(\theta_0 + \theta_x) \operatorname{sgn}(\Re(\theta_0 + \theta_x)), (\theta_0 - \theta_x) \operatorname{sgn}(\Re(\theta_0 - \theta_x))\}$, or $\Sigma_{\beta\delta} = \{(\theta_0 + \theta_x), (\theta_0 - \theta_x), -(\theta_0 + \theta_x), -(\theta_0 - \theta_x)\}$. We also introduce the set $\Omega_{\alpha\gamma} = \Omega_{\alpha\gamma}^1 \cup \Omega_{\alpha\gamma}^2$, where

$$\Omega_{\alpha\gamma}^k := \left\{ \left(\sqrt{2\alpha} + (-)^k \sqrt{2\gamma} \right) \operatorname{sgn} \left(\Re(\sqrt{2\alpha} + (-)^k \sqrt{2\gamma}) \right) \right\}, \quad k = 1, 2 \quad (7)$$

or, in case $|\Re\{\sqrt{2\alpha} + (-)^k \sqrt{2\gamma}\}| < 1$,

$$\Omega_{\alpha\gamma}^k := \left\{ \left(\sqrt{2\alpha} + (-)^k \sqrt{2\gamma} \right), -\left(\sqrt{2\alpha} + (-)^k \sqrt{2\gamma} \right) \right\}. \quad (8)$$

Again, this is $\Omega_{\alpha\gamma} = \{(\theta_\infty - 1 + \theta_1) \operatorname{sgn}(\Re(\theta_\infty - 1 + \theta_1)), (\theta_\infty - 1 - \theta_1) \operatorname{sgn}(\Re(\theta_\infty - 1 - \theta_1))\}$, or $\Omega_{\alpha\gamma} = \{(\theta_\infty - 1 + \theta_1), (\theta_\infty - 1 - \theta_1), -(\theta_\infty - 1 + \theta_1), -(\theta_\infty - 1 - \theta_1)\}$

The behaviors in the table may be classified as follows:

- Complex power behaviors (22), (26), (37) and (41). They are expanded in powers of $x^{n+m\lambda}$, for some $n, m \in \mathbf{Z}$ and $\lambda \in \mathbf{C}$. In this case, $|y(x)|$ may vanish, converge to a constant or diverge when $x \rightarrow 0$.
- Inverse oscillatory behaviors (35) and (39). $y(x)$ oscillates without vanishing when $x \rightarrow 0$, and may have poles in a sector centered at $x = 0$.
- Integer power behaviors, namely Taylor series (30), (31), (45), (46), (27) and (38).
- Power logarithmic behaviors, of type (28), (29), (33) and (43), namely power expansions with coefficients which are polynomials of $\ln x$. $y(x) \rightarrow \text{constant}$ as $x \rightarrow 0$, where the constant may be zero or not.
- Inverse power logarithmic behaviors of type (44) and (48), which can be expanded as asymptotic series of $(\ln x)^{-1}$. $y(x) = O(1/\ln x)$, or $O(1/\ln^2 x)$, as $x \rightarrow 0$.

Except for the case when logarithms appear in the expansions tabulated, the expansions are convergent and define true solutions. The proof is based on the results of [22] and [9]. When logarithms appear, no proof of convergence is known to the author, though one may expect that an extension of the method of [22] and [9] be possible for the power logarithmic expansions.

In the table, σ, ϕ and a denote complex free parameters (integration constants), and ν a real free parameter. The coefficients c_{nm} , d_{nm} and b_n are rational functions of $\sqrt{\alpha}$, $\sqrt{\beta}$, $\sqrt{\gamma}$ and $\sqrt{1-2\delta}$. The coefficients $b_n(a)$ are rational functions of $\sqrt{\alpha}$, $\sqrt{\beta}$, $\sqrt{\gamma}$, $\sqrt{1-2\delta}$ and a . $P_n(\ln x, a)$ are polynomials in $\ln x$ with coefficients which are rational functions of $\sqrt{\alpha}$, $\sqrt{\beta}$, $\sqrt{\gamma}$, $\sqrt{1-2\delta}$ and a . These coefficients can be recursively computed (essentially, by substitution into PVI. See [12]). In the case of (22), the c_{nm} 's depend rationally also on σ . In the case of (35), the c_{nm} 's depend rationally also on ν .

Complex power behaviors		Free Parameters	Other Conditions
(22)	$y(x) = \sum_{n=1}^{\infty} x^n \sum_{m=-n}^n c_{nm} (ax^{\sigma})^m$ Basic solutions	$\sigma, \ a \neq 0$	$0 \leq \Re \sigma < 1, \ \sigma \notin \Sigma_{\beta\delta},$ $c_{11} = 1$
(26)	$y(x) = \sum_{n=1}^{\infty} x^n \sum_{m=0}^n c_{nm} (ax^{\sigma})^m$ Basic solutions	$a \neq 0$	$\Re \sigma > -1, \ \sigma \in \Sigma_{\beta\gamma}^k,$ $\Sigma_{\beta\gamma}^k \cap \mathbf{Z} = \emptyset, \ k = 1, 2.$ $\Sigma_{\beta\gamma}^k$ is (5) or (6). $c_{11} = 1,$ $c_{00} = \frac{\sqrt{-2\beta}}{\sqrt{-2\beta+(-)^k\sqrt{1-2\delta}}}$
(37)	$y(x) = \frac{\sqrt{\alpha+(-)^k\sqrt{\gamma}}}{\sqrt{\alpha}} + \sum_{n=1}^{\infty} x^n \sum_{m=0}^n d_{nm} (ax^{\rho})^m$	a	$\Re \rho > -1, \ \rho + 1 \in \Omega_{\alpha\gamma}^k,$ $\Omega_{\alpha\gamma}^k \cap \mathbf{Z} = \emptyset, \ k = 1, 2.$ $\Omega_{\alpha\gamma}^k$ is (7)
(41)	$y(x) = \frac{1}{a} x^{-\omega} (1 + \sum_{n=1}^{\infty} x^n \sum_{m=0}^n d_{nm} (ax^{\omega})^m)$	a	$\alpha = 0.$ $\omega \notin \mathbf{Z}, \ \Re \omega \geq 0.$ $\omega \in \{\sqrt{2\gamma}, -\sqrt{2\gamma}\}$ if $\gamma < 0,$ otherwise $\omega = \sqrt{2\gamma} \operatorname{sgn}(\Re \sqrt{2\gamma}).$

Inverse Oscillatory Behaviors		Free Param	Other
(35)	$y(x) = \left[\sum_{n=0}^{\infty} x^n \sum_{m=-n-1}^{n+1} c_{nm} (e^{i\phi} x^{2i\nu})^m \right]^{-1}$ $= [A \sin(2\nu \ln x + \phi) + B + O(x)]^{-1}$	$\nu, \ \phi$	$A = -\sqrt{\frac{\alpha}{2\nu^2} + B^2}$ $B = \frac{2\nu^2 + \gamma - \alpha}{4\nu^2},$ $\nu \in \mathbf{R} \setminus \{0\}, \ 2i\nu \notin \Omega_{\alpha\gamma}$
(39)	$y(x) = \left[\frac{\sqrt{\alpha}}{\sqrt{\alpha+(-)^k\sqrt{\gamma}}} + ax^{2i\nu} + \sum_{n=1}^{\infty} x^n \sum_{m=0}^{n+1} c_{n+1,m} (ax^{2i\nu})^m \right]^{-1}$	a	$\nu \in \mathbf{R} \setminus \{0\}, \ 2i\nu \in \Omega_{\alpha\gamma}^k,$ $k = 1, 2.$ $\Omega_{\alpha\gamma}^k \subset i\mathbf{R}$ is (8)

Taylor expansions	Free Par	Other Conditions
(27) $y(x) = \frac{\sqrt{-2\beta}}{\sqrt{-2\beta \pm \sqrt{1-2\delta}}} x + \sum_{n=2}^{\infty} b_n x^n$		$y(x) = 0$ if $\beta = 0$. $\sqrt{-2\beta} \pm \sqrt{1-2\delta} \notin \mathbf{Z}$.
(30) $y(x) = \sum_{n=1}^{ N } b_n x^n + ax^{ N +1} + \sum_{n= N +2}^{\infty} b_n(a)x^n$ (31) $y(x) = ax + a(a-1)\left(\gamma - \alpha - \frac{1}{2}\right)x^2 + \sum_{n=3}^{\infty} b_n(a)x^n, \quad N=0$	a	One of $\sqrt{-2\beta} + \sqrt{1-2\delta}$ or $\sqrt{-2\beta} - \sqrt{1-2\delta} = N \in \mathbf{Z}$ and $2\beta = -N^2$, or $\{\sqrt{2\alpha} + \sqrt{2\gamma}, \sqrt{2\alpha} - \sqrt{2\gamma}\} \cap \{ N - 1, 1 - N \} \neq \emptyset$. (* $2\beta = 2\delta - 1 = 0$ when $N = 0$)
(38) $y(x) = \frac{\sqrt{\alpha} \pm \sqrt{\gamma}}{\sqrt{\alpha}} + \sum_{n=1}^{\infty} b_n x^n$, Basic Taylor		$\alpha \neq 0$, $\sqrt{2\alpha} \pm \sqrt{2\gamma} \notin \mathbf{Z}$
(45) $y(x) = \sum_{n=0}^{ N -1} b_n x^n + ax^{ N } + \sum_{n= N +1}^{\infty} b_n(a)x^n$ (46) $y(x) = a + (1-a)(\delta - \beta)x + \sum_{n=2}^{\infty} b_n(a)x^n, \quad N=0$ Two Basic Taylor solutions when $N = 0, 1$	a	One of $\sqrt{2\alpha} + \sqrt{2\gamma}$ or $\sqrt{2\alpha} - \sqrt{2\gamma} = N \in \mathbf{Z}$ and $2\alpha = N^2$, or $\{\sqrt{-2\beta} + \sqrt{1-2\delta}, \sqrt{-2\beta} - \sqrt{1-2\delta}\} \cap \{ N - 1, 1 - N \} \neq \emptyset$ (* $\alpha = \gamma = 0$ when $N = 0$)

Power logarithmic behaviors		Free Par.	Other Conditions
(28)	$y(x) = \sum_{n=1}^{ N } b_n x^n + \left(a + b_{ N +1} \ln x \right) x^{ N +1} + \sum_{n= N +2}^{\infty} P_n(\ln x; a) x^n$	a	$\sqrt{-2\beta} + \sqrt{1-2\delta} \text{ or } \sqrt{-2\beta} - \sqrt{1-2\delta} = N \in \mathbf{Z}$ $(*2\beta = 2\delta - 1 \text{ if } N = 0)$ and $2\beta^2 \neq -N^2,$ $\sqrt{2\alpha} \pm \sqrt{2\gamma} \neq \pm(N - 1)$
(29)	$y(x) = \left(a \pm \sqrt{-2\beta} \ln x \right) x + \sum_{n=2}^{\infty} P_n(\ln x; a) x^n, \quad N = 0$ Basic solution when $N = 0$		
(33)	$y(x) = \left[\frac{2\beta+1-2\delta}{4} (a + \ln x)^2 + \frac{2\beta}{2\beta+1-2\delta} \right] x + \sum_{n \geq 2}^{\infty} P_n(\ln x; a) x^n$ Basic solution	a	$2\beta \neq 2\delta - 1$
(43)	$y(x) = \sum_{n=0}^{ N -1} b_n x^n + \left(a + b_N \ln x \right) x^{ N } + \sum_{n= N +1}^{\infty} P_n(\ln x; a) x^n$	a	$\sqrt{2\alpha} + \sqrt{2\gamma} \text{ or } \sqrt{2\alpha} - \sqrt{2\gamma} = N \in \mathbf{Z} \setminus \{0\},$ and $2\alpha \neq N^2,$ $\sqrt{-2\beta} \pm \sqrt{1-2\delta} \neq \pm(N - 1)$

Inverse power logarithmic behaviors		Free Parameters	Other Conditions
(44)	$y(x) = \left\{ \left(a \pm \sqrt{2\alpha} \ln x \right) + \sum_{n=1}^{\infty} P_n(\ln x; a) x^n \right\}^{-1}$ $= \pm \frac{1}{\sqrt{2\alpha} \ln x} \left[1 \mp \frac{a}{\sqrt{2\alpha} \ln x} + O\left(\frac{1}{\ln^2 x}\right) \right]$	a	$\alpha = \gamma \neq 0$
(48)	$y(x) = \left\{ \left[\frac{\gamma-\alpha}{2} (a + \ln x)^2 + \frac{\alpha}{\alpha-\gamma} \right] + \sum_{n=1}^{\infty} P_{n+1}(\ln x; a) x^n \right\}^{-1} =$ $= \frac{2}{(\gamma-\alpha) \ln^2 x} \left[1 - \frac{2a}{\ln x} + O\left(\frac{1}{\ln^2 x}\right) \right]$	a	$\alpha \neq \gamma$

3 Table of Parametrization formulas

The critical behaviors are governed by two integration constants, or free parameters, which appear in the explicit expansions at $x = 0$. We give a tabulation of the parametrization formulas of these constants, in terms of monodromy data of the associated system (1). As for their computation is concerned, the reader may see [17] and [2] for (22) and (26), and my papers [10], [11] and [12] for the other cases.

Parametrization formulas are the achievement of the method of monodromy preserving deformations. They can be written at $x = 0, 1, \infty$ for the same monodromy data, namely for the same $y(x)$. This allows to solve the connection problem.

From the table of the critical behaviors, we extract a subset of behaviors which are *basic* (see section 5 for a precise definition). All the other behaviors in the table can be derived from the basic ones, by the action of a birational Okamoto's transformation of section 5. All the behaviors at $x = 1, \infty$ can also be derived from those of the table, by an Okamoto's transformation (12) and (13), as it is explained in section 5. In this sense, all the information we need is contained in the basic behaviors listed below:

◇ Solutions (22) and (26).

◇ The power logarithmic behaviors (29) and (33):

$$\begin{cases} (29) & y(x) = \left(a \pm \sqrt{-2\beta} \ln x\right)x + \sum_{n=2}^{\infty} P_n(\ln x; a)x^n, & 2\beta = 2\delta - 1 \\ (33) & y(x) = \left[\frac{2\beta+1-2\delta}{4}(a + \ln x)^2 + \frac{2\beta}{2\beta+1-2\delta}\right]x + \sum_{n=2}^{\infty} P_n(\ln x; a)x^n, & 2\beta \neq 2\delta - 1 \end{cases}$$

It is enough to compute the parametrization formulas of the integration constants for the basic critical behaviors (22), (26), (29) and (33). The other critical behaviors are obtained from the basic ones by birational transformations, and the parametrization formulas are correspondingly obtained by an action of the birational transformations on the monodromy data. Examples of these actions are formulas (52) and (53) to be introduced later, which allow to pass from $x = 0$ to $x = 1$ and $x = \infty$ respectively, and (54), associated to the birational transformation (14).

We give also the parametrization formulas for the solutions (35) and (39), which are not basic, but are important for their behavior and their applications in quantum cohomology (see for example [13]). We give also the parametrization formulas for the Taylor solutions below:

$$\begin{cases} (46) & y(x) = a + (1-a)(\delta - \beta)x + \sum_{n=2}^{\infty} b_n(a)x^n, & \alpha = \gamma = 0 \\ (47) & y(x) = \pm \frac{1}{\sqrt{2\alpha}} + ax + \sum_{n=2}^{\infty} b_n(a)x^n, & \alpha \neq 0, (\sqrt{2\alpha} \pm \sqrt{2\gamma})^2 = 1, 2\beta = 2\delta - 1 \end{cases}$$

The latter of these is one of the two solutions (45) when $N = 1$. These solutions, together with (38) (which has no free parameter), are *basic Taylor solutions*, which generate all the other Taylor expansions via birational transformations. We will explain in section 5 that (46) and (47) can be obtained from (29).

Let $p_{ij} = \text{Tr}(M_i M_j)$, $i, j = 0, x, 1$.

Case of (22) and (26) . The two integration constants are σ and a . Their parametrization is computed in [17] and [2]) (note that in [17], the last sign in formula (1.8) at the bottom of page 1141 is $\pm\sigma$ instead of $\mp\sigma$):

$$\sigma = \frac{1}{\pi} \arccos \left(\frac{p_{0x}}{2} \right),$$

$$a = \frac{(\theta_x - \theta_0 - \sigma)(\theta_x + \theta_0 - \sigma)(\theta_\infty + \theta_1 - \sigma)}{4\sigma^2(\theta_\infty + \theta_1 + \sigma)\mathbf{F}}$$

where

$$\mathbf{F} := \frac{\Gamma(1+\sigma)^2 \Gamma\left(\frac{1}{2}(\theta_0 + \theta_x - \sigma) + 1\right) \Gamma\left(\frac{1}{2}(\theta_x - \theta_0 - \sigma) + 1\right)}{\Gamma(1-\sigma)^2 \Gamma\left(\frac{1}{2}(\theta_0 + \theta_x + \sigma) + 1\right) \Gamma\left(\frac{1}{2}(\theta_x - \theta_0 + \sigma) + 1\right)} \times$$

$$\times \frac{\Gamma\left(\frac{1}{2}(\theta_\infty + \theta_1 - \sigma) + 1\right) \Gamma\left(\frac{1}{2}(\theta_1 - \theta_\infty - \sigma) + 1\right)}{\Gamma\left(\frac{1}{2}(\theta_\infty + \theta_1 + \sigma) + 1\right) \Gamma\left(\frac{1}{2}(\theta_1 - \theta_\infty + \sigma) + 1\right)} \frac{V}{U},$$

and:

$$\begin{aligned}
U &:= \left[\frac{i}{2} \sin(\pi\sigma) p_{x1} - \cos(\pi\theta_x) \cos(\pi\theta_\infty) - \cos(\pi\theta_0) \cos(\pi\theta_1) \right] e^{i\pi\sigma} + \\
&\quad + \frac{i}{2} \sin(\pi\sigma) p_{01} + \cos(\pi\theta_x) \cos(\pi\theta_1) + \cos(\pi\theta_\infty) \cos(\pi\theta_0) \\
V &:= 4 \sin \frac{\pi}{2} (\theta_0 + \theta_x - \sigma) \sin \frac{\pi}{2} (\theta_0 - \theta_x + \sigma) \sin \frac{\pi}{2} (\theta_\infty + \theta_1 - \sigma) \sin \frac{\pi}{2} (\theta_\infty - \theta_1 + \sigma).
\end{aligned}$$

Note that for (26), one has to take the limit of the above a for σ tending to one of $\pm(\theta_0 + \theta_x)$ or $\pm(\theta_0 - \theta_x)$. The limit exist finite and non zero.

For (22) and (26), we have

$$\text{Tr}(M_0 M_x) \notin (-\infty, -2] \cup \{2\}$$

In particular

$$\text{Tr}(M_0 M_x) > 2 \text{ when } \sigma \in i\mathbf{R}$$

Case of solutions (35) and (39). The constants ν and ϕ of (35) in terms of the monodromy data are computed in my paper [12].

$$2 \cosh(2\pi\nu) = -p_{0x}, \quad \phi = i \ln \left(-\frac{r}{\nu A} \right)$$

A is the coefficient multiplying the sine in (35):

$$A = \frac{1}{8\nu^2} \sqrt{[4\nu^2 + (\theta_\infty - 1 - \theta_1)^2][4\nu^2 + (\theta_\infty - 1 + \theta_1)^2]}$$

and

$$r = \frac{(\theta_\infty - 1 - \theta_1 - 2i\nu)(\theta_\infty - 1 + \theta_1 + 2i\nu)(\theta_0 + \theta_x + 1 + 2i\nu)}{8i\nu(2i\nu - \theta_0 - \theta_x - 1)} \frac{1}{\mathbf{F}^*},$$

where

$$\begin{aligned}
\mathbf{F}^* &:= \frac{\Gamma(1 - 2i\nu)^2 \Gamma\left(\frac{1}{2}(\theta_\infty - 1 + \theta_1 + 2i\nu) + 1\right) \Gamma\left(\frac{1}{2}(\theta_1 + 1 - \theta_\infty + 2i\nu) + 1\right)}{\Gamma(1 + 2i\nu)^2 \Gamma\left(\frac{1}{2}(\theta_\infty - 1 + \theta_1 - 2i\nu) + 1\right) \Gamma\left(\frac{1}{2}(\theta_1 + 1 - \theta_\infty - 2i\nu) + 1\right)} \times \\
&\quad \times \frac{\Gamma\left(\frac{1}{2}(\theta_0 + \theta_x + 1 + 2i\nu) + 1\right) \Gamma\left(\frac{1}{2}(\theta_x - \theta_0 + 1 + 2i\nu)\right)}{\Gamma\left(\frac{1}{2}(\theta_0 + \theta_x + 1 - 2i\nu) + 1\right) \Gamma\left(\frac{1}{2}(\theta_x - \theta_0 + 1 - 2i\nu)\right)} \frac{\mathcal{V}}{\mathcal{U}}
\end{aligned}$$

and:

$$\begin{aligned}
\mathcal{U} &:= e^{2\pi\nu} \left[\frac{1}{2} \sinh(2\pi\nu) p_{1x} + \cos(\pi\theta_x) \cos(\pi\theta_\infty) + \cos(\pi\theta_0) \cos(\pi\theta_1) \right] + \\
&\quad - \frac{1}{2} \sinh(2\pi\nu) p_{01} + \cos(\pi\theta_x) \cos(\pi\theta_1) + \cos(\pi\theta_\infty) \cos(\pi\theta_0)
\end{aligned}$$

$$\mathcal{V} := 4 \cos \frac{\pi}{2} (\theta_0 + \theta_x + 2i\nu) \cos \frac{\pi}{2} (\theta_0 - \theta_x - 2i\nu) \sin \frac{\pi}{2} (\theta_\infty - 1 + \theta_1 + 2i\nu) \sin \frac{\pi}{2} (\theta_\infty - 1 - \theta_1 - 2i\nu).$$

As for the solution (39), we have

$$a = -\frac{r}{2i\nu}$$

where r is the above. Note that in this case $2i\nu$ is equal to one of $\pm(\theta_\infty - 1 + \theta_1)$ or $\pm(\theta_\infty - 1 - \theta_1)$, and the limit of r above exists for these values of $2i\nu$.

For (35) and (39), we have

$$\text{Tr}(M_0 M_x) < -2$$

Case of (47): For the basic Taylor expansion (47), the parametrization is computed in my paper [10]. Note that $\theta_\infty \neq 1$, one of $\theta_0 \pm \theta_x = 0$, and one of $\theta_\infty - 1 \pm \theta_1 = 1$ or -1 . The parameterization is:

$$y(x) = \frac{1}{1 - \theta_\infty} + a x + O(x^2),$$

$$a = \frac{\theta_\infty(2s + \theta_x + 1)}{2(\theta_\infty - 1)}, \quad s = \frac{\theta_x[2 \cos(\pi(\theta_\infty + \theta_x)) - p_{01}]}{2[\cos(\pi(\theta_\infty - \theta_x)) - \cos(\pi(\theta_\infty + \theta_x))]}.$$

In this case

$$\text{Tr}(M_0 M_x) = 2, \quad \langle M_0 \cdot M_x, M_1 \rangle \text{ is reducible}$$

Case of (46): For the basic Taylor expansion (46), the parametrization is computed in my paper [10]. Note that one of $\theta_\infty - 1 \pm \theta_1 = 0$. The parameterization is:

$$y(x) = a + \frac{1-a}{2}(1 + \theta_0^2 - \theta_x^2)x + O(x^2)$$

$$a = \frac{1}{1-s}, \quad s = \frac{p_{01} - 2 \cos(\pi \theta_0)}{4 \cos\left(\frac{\pi \theta_0}{2} - \frac{\pi \theta_x}{2}\right) \cos\left(\frac{\pi \theta_0}{2} + \frac{\pi \theta_x}{2}\right)}$$

This result can also be obtained from (9) below, via (14) and (54). In case of (46) we have:

$$\text{Tr}(M_0 M_x) = -2, \quad \langle M_0 \cdot M_x, M_1 \rangle \text{ is reducible}$$

Case of (29): The free parameter of (29) is computed in my paper [11], for $\theta_\infty \pm \theta_1 \notin 2\mathbf{Z}$. Note that one of $\theta_x \pm \theta_0 = 0$. The parameterization is:

$$a = e^{i\frac{\pi}{2}\theta_0}\theta_0 \left\{ \Psi_E(\theta_0 + 1) + i\pi + \gamma_E - \frac{\pi}{4} \frac{p_{01}}{\sin \pi \theta_0 \sin \frac{\pi}{2}(\theta_\infty + \theta_1) \sin \frac{\pi}{2}(\theta_\infty - \theta_1)} + \right.$$

$$\left. + \frac{\pi}{2} \frac{\cos \pi(\theta_0 + \theta_1)}{\sin \pi \theta_0 \sin \frac{\pi}{2}(\theta_\infty + \theta_1) \sin \frac{\pi}{2}(\theta_\infty - \theta_1)} - \frac{\omega}{2} \frac{[\cos \pi(\theta_0 + \theta_1) - \cos \pi(\theta_0 - \theta_1)]}{\sin \pi \theta_0 \sin \pi \theta_1} \right\}$$

$$\omega = \Psi_E\left(\frac{\theta_\infty}{2} + \frac{\theta_1}{2}\right) - \Psi_E\left(\frac{\theta_1}{2} - \frac{\theta_\infty}{2} + 1\right) + 2\gamma_E$$

Here Ψ_E is the Euler's psi-function, and γ_E is the Euler's gamma constant. In this case

$$\text{Tr}(M_0 M_x) = 2, \quad \langle M_0, M_x \rangle \text{ is reducible}$$

When $\theta_0 \rightarrow 0$, namely $\beta \rightarrow 0$, (29) becomes the Taylor series (31) and

$$a \rightarrow \frac{2 \cos \pi \theta_1 - p_{01}}{4 \sin \frac{\pi}{2}(\theta_\infty + \theta_1) \sin \frac{\pi}{2}(\theta_\infty - \theta_1)} \quad (9)$$

Case (33): The free parameter for (33) is calculated in my paper [11], when $\theta_0, \theta_x, \theta_1, \theta_\infty \notin \mathbf{Z}$. Note that $\theta \pm \theta_x \neq 0$. The parameterization is:

$$a = \frac{C - c + p_{0x} - p_{x1}}{b - B} + \frac{2\theta_x}{\theta_0^2 - \theta_x^2} + 4(\gamma_E - i\pi) +$$

$$+ \Psi_E\left(\frac{\theta_\infty}{2} + \frac{\theta_1}{2}\right) - \Psi_E\left(\frac{\theta_1}{2} - \frac{\theta_\infty}{2} + 1\right) + \Psi_E\left(-\frac{\theta_x}{2} - \frac{\theta_0}{2}\right) + \Psi_E\left(\frac{\theta_x}{2} - \frac{\theta_0}{2} + 1\right)$$

where

$$b = \frac{4}{\pi}(\sin \pi \theta_1 s(\theta_0 - \theta_x)s(\theta_0 + \theta_x) + \sin \pi \theta_0 s(\theta_\infty + \theta_1)s(\theta_\infty - \theta_1))$$

$$c = 2 \cos \pi(\theta_0 - \theta_1)$$

$$B = \frac{1}{2\pi i} \left[2 \cos \pi(\theta_0 + \theta_1) + 4 \cos \pi \theta_x \cos \pi \theta_\infty - 4e^{i\pi \theta_1} \cos \pi \theta_x - 4e^{-i\pi \theta_0} \cos \pi \theta_\infty + 3e^{i\pi(\theta_1 - \theta_0)} - e^{i\pi(\theta_0 - \theta_1)} \right]$$

$$C = 2e^{i\pi \theta_1} \cos \pi \theta_x + 2e^{-i\pi \theta_0} \cos \pi \theta_\infty - 2e^{i\pi(\theta_1 - \theta_0)}$$

and $s(z) := \sin(\pi z/2)$. In this case

$$\text{Tr}(M_0 M_x) = 2$$

4 Completeness of the Table of Critical Behaviors

The monodromy data θ_0 , θ_1 , θ_x and θ_∞ are fixed (up to $\theta_k \mapsto -\theta_k$, $k = 0, x, 1$, and $\theta_\infty \mapsto 2 - \theta_\infty$) by PVI. Consider a triple of 2×2 matrices M_0 , M_x and M_1 such that

$$\text{Det} M_0 = \text{Det} M_1 = \text{Det} M_x = 1, \quad \text{Tr} M_\mu = 2 \cos \pi \theta_\mu, \quad \mu = 0, 1, x, \infty$$

where $M_\infty := M_1 M_x M_0$. These matrices are defined up to conjugation by an invertible matrix. The set of such triples, modulo conjugation, is the set of possible monodromy matrices for the system (1) with the basis of loops ordered as figure 1. We call it, together with θ_0 , θ_1 , θ_x and θ_∞ , the *monodromy data* of Fuchsian systems (1), with the basis of loops ordered as figure 1. For an equivalent definition, see section 2 of [12].

To every branch $y(x)$ of a solution of PVI, a system (1) is associated, and its monodromy group. Therefore, to $y(x)$ a point in the space of monodromy data is associated. Conversely, to a point in the space of monodromy data, either one branch $y(x)$ may be uniquely associated, or an entire family of solutions may be associated. Also, no solution of PVI may correspond, in principle, to some exceptional points in the space of monodromy data. In [11] we proved that a branch $y(x)$ is uniquely associated to a point in the space of monodromy data if and only if:

a*) none of the monodromy matrices M_0 , M_x , M_1 and $M_1 M_x M_0$ (order of figure 1) is the identity.

Let $p_{ij} := \text{Tr}(M_i M_j)$, $j = 0, x, 1$ and $p_\mu := \text{Tr} M_\mu = 2 \cos \pi \theta_\mu$, $\mu = 0, x, 1, \infty$. It is known (see [16]) that p_{0x} , p_{x1} , p_{1x} and the p_μ 's (equivalently, the θ_μ 's up to sign) are good coordinates for the monodromy data when:

b*) the group generated by M_0 , M_x , M_1 is irreducible.

If a*) and b*) hold, we can parameterize the integration constants of the critical behavior of a branch $y(x)$ in terms of the p_{ij} 's and θ_μ 's, as in section 3. See section 8 for more details.

A natural question is if the table of critical behaviors is complete, namely if it includes all the possible critical behaviors. A way to answer is to see if the correspondence $\{y(x)\} \mapsto \{\text{monodromy data}\}$ is surjective. Namely, the table is complete if the image, through this correspondence, is the whole space of monodromy data.

The results of the previous section, supplemented by the application of birational transformations, bring to the following conclusion:

To the set of complex power behaviors and inverse oscillatory behaviors (namely (22), (26), (37), (41), (35) and (39)) the whole subspace of monodromy data such that $\text{Tr}(M_0 M_x) \neq \pm 2$ is associated. The subspace of monodromy data such that $\text{Tr}(M_0 M_x) = \pm 2$ contains the image of all the remaining solutions in the table of critical behaviors (Taylor, power logarithmic and inverse power logarithmic). This image is the union of the following subsets (let $N \in \mathbf{Z}$):

$$\text{Tr}(M_0 M_x) = 2, \quad \theta_0 \pm \theta_x \neq 0 \text{ and } \langle M_0, M_x, M_1 \rangle \text{ contains no reducible subgroup}$$

$$\begin{aligned} \text{Tr}(M_0 M_x) &= 2, \text{ one of } \theta_0 \pm \theta_x = 2N \text{ and } \langle M_0, M_x \rangle \text{ is reducible} \\ \text{Tr}(M_0 M_x) &= 2, \text{ one of } \theta_\infty - 1 \pm \theta_1 = 2N + 1 \text{ and } \langle M_0 \cdot M_x, M_1 \rangle \text{ is reducible} \end{aligned}$$

$$\text{Tr}(M_0 M_x) = -2, \quad \theta_\infty - 1 \pm \theta_1 \neq 0 \text{ and } \langle M_0, M_x, M_1 \rangle \text{ contains no reducible subgroup}$$

$$\begin{aligned} \text{Tr}(M_0 M_x) &= -2, \text{ one of } \theta_0 \pm \theta_x = 2N + 1 \text{ and } \langle M_0, M_x \rangle \text{ is reducible} \\ \text{Tr}(M_0 M_x) &= -2, \text{ one of } \theta_\infty - 1 \pm \theta_1 = 2N \text{ and } \langle M_0 \cdot M_x, M_1 \rangle \text{ is reducible} \end{aligned}$$

In this sense, the table of the critical behaviors is essentially complete.

To conclude this section, we give more details about the above characterization of $\{y(x)\} \mapsto \{\text{monodromy data}\}$.

- To the space of solutions (22) and (26), and their images through the birational transformation (14) (to be introduced in section 5), namely the solutions (37), (41), (35) and (39), there corresponds the subspace of monodromy data such that

$$\text{Tr}(M_0 M_x) \neq \pm 2$$

- To the space of the power logarithmic solutions (28) and (29), there corresponds the subspace of monodromy data satisfying

$$\text{Tr}(M_0 M_x) = 2 \cos \pi N = \pm 2, \text{ one of } \theta_0 \pm \theta_x = N, \quad \langle M_0, M_x \rangle \text{ is reducible.}$$

Taylor solutions (30) and (31) are included in this case if moreover

$$\theta_0 = \pm N, \text{ or one of } \theta_\infty - 1 \pm \theta_1 = \pm(|N| - 1)$$

- To the space of the power logarithmic solutions (33) there corresponds the subspace of monodromy data such that

$$\text{Tr}(M_0 M_x) = 2, \quad \theta_0 \pm \theta_x \neq 0, \quad \langle M_0, M_x, M_1 \rangle \text{ has no reducible subgroup.}$$

- To the space of the power logarithmic solutions (43), there corresponds the subspace of monodromy data satisfying

$$\text{Tr}(M_0 M_x) = -2 \cos \pi N = \pm 2, \quad N \neq 0, \text{ one of } \theta_\infty - 1 \pm \theta_1 = N, \quad \langle M_0 \cdot M_x, M_1 \rangle \text{ is reducible.}$$

The Taylor expansions (45) are contained in this case, when also

$$\theta_\infty - 1 = \pm N, \text{ or one of } \theta_0 \pm \theta_x = \pm(|N| - 1)$$

- To the space of the Taylor solutions (46) there corresponds the subspace of monodromy data satisfying

$$\text{Tr}(M_0 M_x) = -2, \quad \theta_\infty - 1 = \theta_1 = 0, \quad \langle M_0 \cdot M_x, M_1 \rangle \text{ is reducible.}$$

- To the space of the inverse power logarithmic solutions (44) there corresponds the subspace of monodromy data such that

$$\text{Tr}(M_0 M_x) = -2, \text{ one of } \theta_\infty - 1 \pm \theta_1 = 0, \quad \langle M_0 \cdot M_x, M_1 \rangle \text{ is reducible.}$$

- To the space of the inverse power logarithmic solutions (48) there corresponds the subspace of the monodromy data such that

$$\text{Tr}(M_0 M_x) = -2, \quad \theta_\infty - 1 \pm \theta_1 \neq 0, \quad \langle M_0, M_x, M_1 \rangle \text{ has no reducible subgroup.}$$

- When

$$\langle M_0, M_x, M_1 \rangle \text{ is reducible}$$

then, according to [14], all the corresponding solutions of (PVI) are equivalent by birational canonical transformations to the following one-parameter family of solutions, with $\theta_\infty + \theta_1 + \theta_0 + \theta_x = 0$:

$$y(x) = \frac{\theta_1 + \theta_\infty - 1 + x(1 + \theta_x)}{\theta_\infty - 1} - \frac{1}{\theta_\infty - 1} \frac{x(1 - x)}{u(x; a)} \frac{du(x; a)}{dx}, \quad (10)$$

where $u(x; a) = u_1(x) + au_2(x)$; $a \in \mathbf{C}$, $u_1(x)$ and $u_2(x)$ are linear independent solutions of the hypergeometric equation:

$$x(1 - x) \frac{d^2 u}{dx^2} + \{[2 - (\theta_\infty + \theta_1)] - (4 - \theta_\infty + \theta_x)x\} \frac{du}{dx} - (2 - \theta_\infty)(1 + \theta_x)u = 0$$

Their behavior is again power logarithmic, Taylor or (37), but the parametrization in terms of p_{0x} , p_{x1} , p_{01} does not apply. The rational solutions fall into the above case, see [20].

5 Okamoto's Birational Transformations and Basic Solutions

Birational transformations are *symmetries* of (PVI), namely invertible transformations:

$$y'(x) = \frac{P\left(x, y(x), \frac{dy(x)}{dx}\right)}{Q\left(x, y(x), \frac{dy(x)}{dx}\right)}, \quad x' = \frac{p(x)}{q(x)}, \quad (\theta_0, \theta_x, \theta_1, \theta_\infty) \mapsto (\theta'_0, \theta'_x, \theta'_1, \theta'_\infty) \quad (11)$$

such that $y(x)$ satisfies (PVI) with coefficients $\theta_0, \theta_x, \theta_1, \theta_\infty$ and variable x , if and only if $y'(x')$ satisfies (PVI) with coefficients $\theta'_0, \theta'_x, \theta'_1, \theta'_\infty$ and variable x' . The functions P, Q are polynomials; p, q are linear; the transformation of the θ_μ 's is an element of a linear representation of one of the following groups.

* Permutation group:

$$\theta'_0 = \theta_1, \quad \theta'_x = \theta_x, \quad \theta'_1 = \theta_0, \quad \theta'_\infty = \theta_\infty; \quad y'(x') = 1 - y(x), \quad x' = 1 - x. \quad (12)$$

$$\theta'_x = \theta_1, \quad \theta'_1 = \theta_x; \quad \theta'_0 = \theta_0, \quad \theta'_\infty = \theta_\infty; \quad y'(x') = \frac{1}{x}y(x), \quad x' = \frac{1}{x}. \quad (13)$$

$$\theta'_0 = \theta_\infty - 1, \quad \theta'_x = \theta_1, \quad \theta'_1 = \theta_x, \quad \theta'_\infty = \theta_0 + 1; \quad y'(x') = \frac{x}{y(x)}, \quad x = x'. \quad (14)$$

In terms of the coefficients of PVI, the transformation (14) is $\alpha' = -\beta$, $\beta' = -\alpha$, $\gamma' = \frac{1}{2} - \delta$, $\delta' = \frac{1}{2} - \gamma$.

* Weyl Group of the root system D_4 :

w_1 :

$$\theta'_1 = -\theta_1; \quad \theta'_0 = \theta_0, \quad \theta'_x = \theta_x, \quad \theta'_\infty = \theta_\infty.$$

w_2 :

$$\theta'_0 = \frac{\theta_0 + \theta_1 + \theta_x + \theta_\infty}{2} - 1, \quad \theta'_1 = \frac{\theta_0 + \theta_1 - \theta_x - \theta_\infty}{2} + 1,$$

$$\theta'_x = \frac{\theta_0 - \theta_1 + \theta_x - \theta_\infty}{2} + 1, \quad \theta'_\infty = \frac{\theta_0 - \theta_1 - \theta_x + \theta_\infty}{2} + 1$$

w_3 :

$$\theta'_\infty = 2 - \theta_\infty; \quad \theta'_0 = \theta_0, \quad \theta'_x = \theta_x, \quad \theta'_1 = \theta_1.$$

w_4 :

$$\theta'_\infty = 2 - \theta_\infty; \quad \theta'_x = 2 - \theta_x; \quad \theta'_0 = \theta_0, \quad \theta'_1 = \theta_1.$$

The variable $x' = x$, but $y'(x)$ is quite complicated and will not be given here.

* Shift $l_j : v = (v_1, v_2, v_3, v_4) \mapsto v + e_j$, $j = 1, 2, 3, 4$, where $e_1 = (1, 0, 0, 0)$, ..., $e_4 = (0, 0, 0, 1)$:

$$l_1 : \quad \theta'_0 = \theta_0 + 1, \quad \theta'_1 = \theta_1 + 1; \quad \theta'_x = \theta_x, \quad \theta'_\infty = \theta_\infty.$$

$$l_2 : \quad \theta'_0 = \theta_0 + 1, \quad \theta'_1 = \theta_1 - 1; \quad \theta'_x = \theta_x, \quad \theta'_\infty = \theta_\infty.$$

$$l_3 : \quad \theta'_x = \theta_x + 1, \quad \theta'_\infty = \theta_\infty + 1; \quad \theta'_0 = \theta_0, \quad \theta'_1 = \theta_1.$$

$$l_4 : \quad \theta'_x = \theta_x + 1, \quad \theta'_\infty = \theta_\infty - 1; \quad \theta'_0 = \theta_0, \quad \theta'_1 = \theta_1.$$

The variable $x' = x$, but $y'(x)$ is quite complicated and will not be given here.

Transformations w_1, w_2, w_3, w_4 and l_3 are sufficient to generate the other l_j 's, and they give a representation of the affine Weyl group of D_4 . The transformations of $y(x)$ corresponding to w_1, w_2, w_3 and w_4 are obtained from formula (2.10) at page 355 of [21], while the transformation corresponding to l_3 is obtained from formulas (2.5), (2.6) at page 354 of [21], provided that h is substituted with h^+ of (1.13), page 352.

The critical behaviors of $y(x)$ when $x \rightarrow 1$ and $x \rightarrow \infty$ are obtained from the behaviors at $x = 0$. Actually, the transformation (12) exchanges the values 0 and 1 of the independent variable. Therefore, when $x \rightarrow 0$ then $x' \rightarrow 1$. We obtain the behavior of $y'(x')$ at $x' = 1$ from that of $y(x)$ at $x = 0$. The transformation (13) exchanges the values x and 1 of the independent variable. Therefore, if $x \rightarrow 0$, $x' \rightarrow \infty$ and we obtain the behavior of $y'(x')$ from that of $y(x)$.

The symmetry (14) allows to construct new behaviors at $x = 0$ from known ones. An example are the inverse oscillatory behaviors or the inverse power logarithmic behaviors. See section 7.

Definition: A set of *basic solutions* is a set of solutions that can generate all the other solutions of the equation via a birational transformation. In the same way, we call basic expansions those expansions that can generate, via a birational transformation, all the expansions which formally satisfy the equation.

Birational transformations allow us to obtain all the solutions given in the table, form the set of basic solutions (22), (26), (29) and (33).

In order to see this, first observe that the complex power behaviors, the Taylor expansions and the power logarithmic expansions given in the table are the only allowed forms of these kinds. This can be proved by direct substitution into PVI.

As a second step, observe that the complex power behaviors (37) and (41), and the inverse oscillatory behaviors are obtained from (22) and (26) via (14). The inverse power logarithmic behaviors and the power logarithmic behaviors (43) are obtained from (28), (29) and (33) via (14). Details are explained in section 7.

The third step is to prove that one can obtain all solutions (28) from (29). This is done by a birational transformation belonging to the affine Weil group of D_4 . This is because these transformations can be used to change $\theta_0 \pm \theta_x$ into $\theta_0 \pm \theta_x + N$, $N \in \mathbf{Z}$. It is a formidable task to obtain $y(x)$ by repeated applications of birational transformations, but one can avoid this by observing that a transformation (11) applied to a power logarithmic behavior necessarily produces a power logarithmic behavior or an inverse power logarithmic behavior. Then, the conclusion comes from the fact that the expansions in

the table exhaust all possible forms of such behaviors, being relations like $\theta_\mu \pm \theta_\nu \in \mathbf{Z}$, or $\notin \mathbf{Z}$, preserved by birational transformations of the affine D_4 .

Finally, Taylor expansions can all be obtained from (29). For example, when we apply the birational transformation l_3 (or l_3^{-1}) to (29), we obtain the case $N = 1$ of solution (28), namely:

$$y(x) = \pm \sqrt{-2\beta} x + \left(a - \frac{1}{2} \sqrt{-2\beta} (\sqrt{-2\beta} \mp 1) (\alpha - \gamma) \ln x \right) x^2 + \sum_{n=2}^{\infty} x^n \sum_{m=0}^{n-1} b_{nm}(a) \ln^m x, \quad (15)$$

$$\sqrt{-2\beta} + \sqrt{1-2\delta} = \pm 1 \text{ or } \sqrt{-2\beta} - \sqrt{1-2\delta} = \pm 1$$

Then, let $\beta = 0$ in (29) and $\alpha = \gamma$ in (15). We obtain respectively the two basic Taylor solutions:

$$y(x) = a + \frac{a(a-1)}{2} (2\gamma - 2\alpha - 1) x^2 + \sum_{n=3}^{\infty} b_n(a) x^n, \quad \text{this is (31) in the table}$$

$$y(x) = \pm \sqrt{-2\beta} x + ax^2 + \sum_{n=3}^{\infty} b_n(a) x^n, \quad \text{this is (32), section 7}$$

These are transformed by (14) into (46) and (47), which may also be used as basic Taylor solutions. Then, birational transformations belonging to the affine Weil group of D_4 applied the basic Taylor give all the other Taylor solutions at $x = 0$. $\square$

6 How to obtain the Critical Behaviors: Matching Procedure

The matching procedure, supplemented by the symmetries of PVI, gives the critical behaviors at $x \rightarrow 0$, in the table. Originally, such procedure was suggested by Its and Novokshenov in [15], for the second and third Painlevé equations. This approach was further developed and used by Kapaev, Kitaev, Andreev, and Vartanian (see for example [1]). It was implemented in my paper [10] for the case of PVI. We review here the method of [10].

Let us divide the λ -plane into two domains. The “outside” domain is defined for $|\lambda| \geq |x|^{\delta_{OUT}}$, $\delta_{OUT} > 0$. In this domain, (1) can be approximated by:

$$\frac{d\Psi_{OUT}}{d\lambda} = \left[\frac{A_0 + A_x}{\lambda} + \frac{x A_x}{\lambda^2} \sum_{n=0}^{N_{OUT}} \left(\frac{x}{\lambda} \right)^n + \frac{A_1}{\lambda - 1} \right] \Psi_{OUT}, \quad (16)$$

The “inside” domain is defined for λ comparable with x , namely $|\lambda| \leq |x|^{\delta_{IN}}$, $\delta_{IN} > 0$, and (1) can be approximated by:

$$\frac{d\Psi_{IN}}{d\lambda} = \left[\frac{A_0}{\lambda} + \frac{A_x}{\lambda - x} - A_1 \sum_{n=0}^{N_{IN}} \lambda^n \right] \Psi_{IN}, \quad (17)$$

where N_{IN} , N_{OUT} are suitable integers.

The leading term of $y(x)$ is obtained as a result of two facts: a) Systems (16) and (17) are isomonodromic. b) Two fundamental matrix solutions $\Psi_{OUT}(\lambda, x)$, $\Psi_{IN}(\lambda, x)$ must match in the region of overlap, provided this is not empty:

$$\Psi_{OUT}(\lambda, x) \sim \Psi_{IN}(\lambda, x), \quad |x|^{\delta_{OUT}} \leq |\lambda| \leq |x|^{\delta_{IN}}, \quad x \rightarrow 0 \quad (18)$$

This relation is to be intended in the sense that the leading terms of the local behavior of Ψ_{OUT} and Ψ_{IN} for $x \rightarrow 0$ must be equal. It also requires that $\delta_{IN} \leq \delta_{OUT}$.

The simplest reduction is a Fuchsian systems, for $N_{IN} = N_{OUT} = 0$:

$$\frac{d\Psi_{OUT}}{d\lambda} = \left[\frac{A_0 + A_x}{\lambda} + \frac{A_1}{\lambda - 1} \right] \Psi_{OUT}, \quad (19)$$

$$\frac{d\Psi_{IN}}{d\lambda} = \left[\frac{A_0}{\lambda} + \frac{A_x}{\lambda - x} \right] \Psi_{IN}. \quad (20)$$

Practically, we match a fundamental solution of (16) for $\lambda \rightarrow 0$, with a fundamental solution of the system obtained from (17) by the change of variables $\mu := \lambda/x$, namely with a solution of:

$$\frac{d\Psi_{IN}}{d\mu} = \left[\frac{A_0}{\mu} + \frac{A_x}{\mu - 1} - xA_1 \sum_{n=0}^{N_{IN}} x^n \mu^n \right] \Psi_{IN}, \quad \mu := \frac{\lambda}{x}. \quad (21)$$

With the substitution $\lambda = \frac{1}{\nu}$ in the system OUT, we can finally write a form for the matching which is well suitable in case that N_{OUT} and/or N_{IN} is greater than zero:

$$\begin{aligned} \frac{d\Psi_{OUT}}{d\nu} &= \left[-xA_x \sum_{n=0}^{N_{OUT}} (x\nu)^n - \frac{A_0 + A_x}{\nu} + \frac{A_1}{\nu(\nu - 1)} \right] \Psi_{OUT} \\ \frac{d\Psi_{IN}}{d\mu} &= \left[-xA_1 \sum_{n=0}^{N_{IN}} (x\mu)^n + \frac{A_0 + A_x}{\mu} + \frac{A_x}{\mu(\mu - 1)} \right] \Psi_{IN} \end{aligned}$$

The matching being

$$\Psi_{OUT}(\nu, x) \sim \Psi_{IN}(\mu, x), \quad \nu, \mu \rightarrow \infty$$

$$|x|^{-\delta_{IN}} \leq |\nu| \leq |x|^{-\delta_{OUT}}, \quad |x|^{\delta_{OUT}-1} \leq |\mu| \leq |x|^{\delta_{IN}-1}.$$

The above matching produces the leading terms of $A_0(x)$, $A_1(x)$ and $A_x(x)$ as $x \rightarrow 0$, therefore it produces the leading term(s) of $y(x)$ in (2).

7 Critical Behaviors when $x \rightarrow 0$. Description of the table

We explain in detail the behaviors in the table. The behavior of case **1)** below was first obtained by Jimbo in its original and fundamental paper [17]. The matching procedure, together with the Okamoto's transformations, allows to obtain all the other behaviors.

In the paper [6] (and in the long series of papers of the authors of [6] which precedes [6]) the method of power geometry (based on Newton polygons) is used, in order to obtain all the formal expansions that satisfy PVI. They are classified in a way different from that we use here. In the course of the discussion below, we compare step by step the results obtained in the framework of the isomonodromy deformations method and the classification of [6], and show that they coincide. See the paragraphs denoted by an asterisk (*).

1) Two-complex parameter $(c_{11}, \sigma \in \mathbf{C})$ solutions [17],[9],[10],[11], [12], [2]:

$$\boxed{y(x) = \sum_{n=1}^{\infty} x^n \sum_{m=-n}^n c_{nm} x^{m\sigma}, \quad -1 \leq \Re\sigma < 1, \quad c_{11} \neq 0} \quad (22)$$

Note that the family is invariant for $\sigma \mapsto -\sigma$, therefore we can restrict to the range

$$\boxed{0 \leq \Re\sigma < 1}$$

The c_{nm} 's are rational functions of c_{11} and σ , and algebraic functions of the coefficients of PVI. Note that in the table we have used a parameter a and put $c_{11} = 1$. Therefore, here, we have $c_{11} = a$ of the table. If σ does not belong to the set (6), then

$$c_{1,-1} = \frac{\left[(\sqrt{-2\beta} - \sqrt{1-2\delta})^2 - \sigma^2\right] \left[(\sqrt{-2\beta} + \sqrt{1-2\delta})^2 - \sigma^2\right]}{16\sigma^4} \frac{1}{c_{11}} \quad (23)$$

and $c_{nm} = q_{nm}(\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}; \sigma) \times (c_{11})^m$, where q_{nm} are rational functions of their arguments. They are determined by the procedure of [12], section 7. If $\Re\sigma > 0$, the dominant term is

$$y(x) = c_{1,-1} x^{1-\sigma} \left(1 + O(x^\sigma, x^{1-\sigma})\right)$$

On the other hand, if $\sigma = -2i\nu$, $\nu \in \mathbf{R}$, solution (22) becomes a three-real parameter solution

$$y(x) = x \left[A \sin(2\nu \ln x + \phi) + B + \sum_{n=1}^{\infty} x^n \sum_{m=-n-1}^{n+1} c_{nm} x^{m\sigma} \right], \quad (24)$$

$$B^2 - A^2 = \frac{\beta}{2\nu^2}, \quad B = \frac{4\nu^2 + 2\beta + 1 - 2\delta}{8\nu^2},$$

$$c_{nm} = \tilde{q}_{nm}(\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}; \nu) \times e^{im\phi}$$

where $\tilde{q}_{nm}$ are rational functions of their arguments, and the role of c_{11} as integration constant is played by ϕ (which is just the log of c_{11}).

Remark: If σ does not belong to the set (6) and $\Re\sigma > 0$, we can also write (22) as

$$\begin{aligned} y(x) &= \sum_{n=0}^{\infty} x^{2n+1} \sum_{m=-n}^{\infty} a_{nm} x^{m\rho} + \sum_{n=1}^{\infty} x^{2n} \sum_{m=-n}^{\infty} a_{nm} x^{m\rho}, \quad \rho = 1 - \sigma, \quad a_{nm} \in \mathbf{C} \\ &= v_\rho x^\rho \left(1 + \sum_{n=1}^{\infty} x^{n\rho} + \sum_{m=1}^{\infty} \sum_{n=n_0}^{\infty} v_{nm} x^{m(1-\rho)+n\rho} \right), \quad v_\rho, v_{nm} \in \mathbf{C}, \quad v_\rho \neq 0 \end{aligned} \quad (25)$$

where $n_0 = (m-1)/2$ if m is odd, $n_0 = m/2 - 1$ if m is even.

The leading term of the expansions (22) was obtained first by Jimbo in [17]. It can also be obtained with the matching procedure of [10], making use of IN and OUT Fuchsian reductions of the system (1). The full expansion (22) can also be obtained by local analysis, with the method of [22], [9]. This method proves also the convergence of the expansions, which therefore define true solutions of PVI, which are *basic solutions*. In [12] a recursive procedure is given to compute the coefficients c_{nm} .

(*) The class of the solutions (22) when $\Re\sigma \neq 0$ is the class $\mathcal{A}_0$ of [6]. In [6] the $\mathcal{A}_0$ -solutions are written as (25), though the range of n, m is looser, namely $n + m > 0$, $n, m \geq 0$.

(*) The image of $(\mathcal{B}_0^+ \cup \mathcal{B}_0^-) \cup (\mathcal{B}_7^+ \cup \mathcal{B}_7^-)$ of [6] through the symmetry (14) is in the sub class of solutions (24).

◇-----◇

2) Suppose that PVI is such that $\boxed{\sqrt{-2\beta} + \sqrt{1-2\delta} \notin \mathbf{Z}}$ or $\boxed{\sqrt{-2\beta} - \sqrt{1-2\delta} \notin \mathbf{Z}}$ (namely, $\theta_0 + \theta_x$ or $\theta_0 - \theta_x \notin \mathbf{Z}$). In this case, PVI admits a one-complex parameter ($c_{11} \in \mathbf{C}$) family of solutions (see [10] and [12])

$$y(x) = \sum_{n=1}^{\infty} x^n \sum_{m=0}^n c_{nm} x^{m\sigma} = \sum_{k=0}^{\infty} y_k(x) (c_{11} x^\sigma)^k, \quad \sigma \in \Sigma_{\beta\delta}, \quad \Re\sigma > -1 \quad (26)$$

The expansion is convergent, as it follows from the local analysis of [22] and [9], so it defines true solutions of PVI, which are basic. The coefficient c_{11} is the free parameter, $c_{11} = a$ of the table. The c_{nm} are functions $q_{nm}(\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}) \times (c_{11})^m$, like in the case of (22). In particular $c_{10} = \frac{\sqrt{-2\beta}}{\sqrt{-2\beta \pm \sqrt{1-2\delta}}}$. There are two possible solutions, corresponding to the signs $\pm$ above, namely to the fact that $\sigma \in \Sigma_{\beta, \delta}^k$, $k = 1, 2$. We stressed that σ is not a free integration constant. The functions $y_k(x)$ are convergent Taylor expansions of the form

$$y_0(x) = \frac{\sqrt{-2\beta}}{\sqrt{-2\beta + (-)^k \sqrt{1-2\delta}}} x + \sum_{n=2}^{\infty} q_{n0}(\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}) x^n, \quad y_0(x) = 0 \text{ if } \beta = 0 \quad (27)$$

$$y_l(x) = O(x^l).$$

If $-1 < \Re \sigma < 1$, the leading terms of $y(x)$ are

$$y(x) = \frac{\sqrt{-2\beta}}{\sqrt{-2\beta \pm \sqrt{1-2\delta}}} x + c_{11} x^{1+\sigma} + \dots$$

If $\beta = 0$, the leading term is always $c_{11} x^{1+\sigma}$ and

$$y(x) = c_{11} x^{1+\sigma} + \sum_{n=2}^{\infty} x^n \sum_{m=1}^n c_{nm} x^{m\sigma}$$

The solution (26) reduces to a Taylor series solution without free parameters, when $c_{11} = 0$, namely

$$y(x) \equiv y_0(x)$$

The method of local analysis of [22] and [9] allows to extend the result of [17] in the case $\Re \sigma \geq 1$, $\sigma \in \Sigma_{\beta, \delta}$, and this precisely yields existence and convergence of solutions (26). Note that when $-1 < \Re \sigma < 1$, they are a subclass of (22). Actually, the coefficients of $x^{-|m|\sigma}$ in (22) vanish when σ belongs to the set (6) (see formula (23)) provided that $\sqrt{-2\beta} \pm \sqrt{1-2\delta} \notin \mathbf{Z}$ (otherwise coefficients diverge). Also note that this happens when we choose $c_{11} \neq 0$ as integration constant. One can choose $c_{1,-1}$ instead, and express c_{11} from (23). In this case, again the coefficients of $x^{|m|\sigma}$ vanish for σ in the set (6) and we again obtain (26) with $\sigma \mapsto -\sigma$, $\Re(-\sigma) > -1$.

(*) The symmetry (14) transforms the set $(\mathcal{B}_1(\text{non log}) \cup \mathcal{B}_2(\text{non log})) \cup (\mathcal{B}_1^+ \cup \mathcal{B}_1^-) \cup (\mathcal{B}_2^+ \cup \mathcal{B}_2^-) \cup \mathcal{B}_6 \cup (\mathcal{B}_6^+ \cup \mathcal{B}_6^-)$ of [6] into the solutions (26) with $\beta \neq 0$. It transforms $\mathcal{C}_0^\infty \cup \mathcal{B}_8 \cup \mathcal{B}_9$ of [6] into the solutions (26) with $\beta = 0$. This will be clear after reading **2.1** below. [Here we use $\mathcal{B}_i(\text{non log})$ to denote the subclass of $\mathcal{B}_i$ such that the coefficients of powers of x are constants, and the power of x have no integer exponent (see **2.1** below). We distinguish it from the subclass $\mathcal{B}_i(\text{log})$ whose coefficients are polynomials of $\ln x$, and the powers of x have integer exponents (for some value of the coefficients of PVI the coefficients may be constants, and the solutions become Taylor series. See **3.1** below). $\mathcal{B}_i = \mathcal{B}_i(\text{non log}) \cup \mathcal{B}_i(\text{log})$.]

We conclude that the class $\mathcal{A}_0 \cup (\mathcal{B}_0^+ \cup \mathcal{B}_0^-) \cup (\mathcal{B}_1(\text{non log}) \cup \mathcal{B}_2(\text{non log})) \cup (\mathcal{B}_1^+ \cup \mathcal{B}_1^-) \cup (\mathcal{B}_2^+ \cup \mathcal{B}_2^-) \cup \mathcal{B}_6 \cup (\mathcal{B}_6^+ \cup \mathcal{B}_6^-) \cup (\mathcal{B}_7^+ \cup \mathcal{B}_7^-) \cup \mathcal{B}_8 \cup \mathcal{B}_9 \cup \mathcal{C}_0^\infty$ of [6] is contained in the union of **1**) $\cup$ **2**) $\cup$ {the image of **1**) $\cup$ **2**) through the symmetry (14)}.

◇

3) If $\boxed{\sqrt{-2\beta} + \sqrt{1-2\delta} = N}$ or $\boxed{\sqrt{-2\beta} - \sqrt{1-2\delta} = N}$ (namely, $\theta_0 + \theta_x = N$ or $\theta_0 - \theta_x = N$), $N \in \mathbf{Z}$, then solution (26) is no longer defined, because some coefficients diverge. We have instead solutions formally like (26), where σ is the integer N and the divergent coefficients c_{nm} are replaced by

polynomials of $\ln x$. Namely, from the results of [10],[11], we see that there exist one-complex parameter ($a \in \mathbf{C}$) solutions

$$\boxed{y(x) = \sum_{n=1}^{|N|} b_n x^n + \left(a + b_{|N|+1} \ln x\right) x^{|N|+1} + \sum_{n=|N|+2}^{\infty} P_n(\ln x; a) x^n, \quad N \neq 0} \quad (28)$$

$$\boxed{y(x) = \left(a \pm \sqrt{-2\beta} \ln x\right) x + \sum_{n=2}^{\infty} P_n(\ln x; a) x^n, \quad \begin{matrix} N = 0 \\ 2\beta = 2\delta - 1 \end{matrix}} \quad (29)$$

where b_n are rational functions of $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}$, and $\sqrt{1-2\delta}$; a is the free complex parameter; $P_n(\ln x; a)$ are polynomials of $\ln x$ of degree $n-|N|$, with coefficients that are rational functions of $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}$ and a . Solution (29) is a *basic solution*. In particular, the above (28) and (29) reduce to convergent Taylor series (studied in [10] and [19]) if either $\{\sqrt{2\alpha} + \sqrt{2\gamma}, \sqrt{2\alpha} - \sqrt{2\gamma}\} \cap \{|N| - 1, 1 - |N|\} \neq \emptyset$, namely $\{\theta_{\infty} + \theta_1 - 1, \theta_{\infty} - \theta_1 - 1\} \cap \{|N| - 1, 1 - |N|\} \neq \emptyset$, or if $2\beta = -N^2$, namely $\theta_0 \in \{N, -N\}$. The series, starting with x -term, are

$$\boxed{y(x) = \sum_{n=1}^{|N|} b_n x^n + a x^{|N|+1} + \sum_{n=|N|+2}^{\infty} b_n(a) x^n, \quad N \neq 0} \quad (30)$$

$$\boxed{y(x) = y(x) = a x + \frac{a(a-1)}{2} (2\gamma - 2\alpha - 1) x^2 + \sum_{n=3}^{\infty} b_n(a) x^n, \quad \begin{matrix} N = 0 \\ 2\beta = 2\delta - 1 = 0 \end{matrix}} \quad (31)$$

Here $b_n(a)$ are rational functions of $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}$ and a . In (31) we have $\beta = 1 - 2\delta = 0$, namely $\theta_x = \theta_0 = 0$. It is the limit of (29) for $\beta \rightarrow 0$.

In [10], the solution (29) is obtained with the matching procedure, making use of Fuchsian IN and OUT systems as reductions of (1). Convergence of (28) and (29) is not proved, but we expect it, the proof being possibly based on an implementation of the method of [22] and [9]. All the coefficients can be recursively determined by direct substitution into PVI.

The Taylor series are obtained by non-Fuchsian IN and OUT systems as reductions of (1). In [10], three basic Taylor series are obtained in this way, with non vanishing constant term. They will be given later, see the solutions (38), (46) and (47). The symmetry (14) transforms (46) and (47) respectively into (31) and the following:

$$y(x) = \sqrt{-2\beta} x + a x^2 + \sum_{n=3}^{\infty} b_n(a; \sqrt{\alpha}, \sqrt{\beta}) x^n, \quad \beta \neq 0, \quad (\sqrt{-2\beta} \pm \sqrt{1-2\delta})^2 = 1, \quad \alpha = \gamma \quad (32)$$

The last solution is included in (30) for $N = 1$. Taylor expansions are also computed in [19], and their convergence is there motivated.

Solutions (31), (32) and (27) are *basic Taylor solutions*, namely all the other Taylor expanded solutions at $x = 0$ can be obtained from them by birational transformation. For example, when $N = 1$, solutions (30) contain two nonequivalent families, namely the family (32) and the following family of solutions:

$$y(x) = x + a x + \frac{1}{2} a(\gamma - \alpha) x^3 + O(x^4), \quad \delta = -\beta = \frac{1}{2}$$

The latter is equivalent to (31) via birational transformation, while (32) is not.

(*) The symmetry (14) transforms the class $\mathcal{B}_1(\log) \cup \mathcal{B}_2(\log)$ of [6] into (28), and $\mathcal{B}_4 \cup \mathcal{B}_5$ into (29).

◇

4) If $\boxed{2\beta \neq 2\delta - 1}$ (namely, $\theta_0 \pm \theta_x \neq 0$), besides log-solutions (28), there are one-complex parameter ($a \in \mathbf{C}$) logarithmic solutions (see [17], [10] and [11]), which are also *a basic solutions*:

$$y(x) = \left[\frac{2\beta+1-2\delta}{4}(a + \ln x)^2 + \frac{2\beta}{2\beta+1-2\delta} \right] x + \sum_{n \geq 2}^\infty P_n(\ln x; a)x^n, \quad a \in \mathbf{C}. \quad (33)$$

The above solutions are obtained in [10] or [11] by the matching procedure, making use of IN and OUT Fuchsian reductions of system (1). As far as convergence is concerned, no proof is published, though we expect that it is convergent and an implementation of the method of [22] should allow to prove this. All the coefficients can be recursively determined by direct substitution into PVI.

(*) The symmetry (14) transforms the class $\mathcal{B}_3$ of [6] into (33).

◇

◇=====◇

In the following, we apply the symmetry (14) to the solutions **1)-4)**, and we obtain new solutions. In all which follows, it is understood that wherever α, β, γ and δ appear in formulas related to **1)-4)** (for example, the coefficients of the expansions), we must substitute $(\alpha, \beta, \gamma, \delta) \mapsto (-\beta, -\alpha, \frac{1}{2} - \delta, \frac{1}{2} - \gamma)$.

1.1a) When σ does not belong to the set (6) and $\Re \sigma > 0$, then from (22) we obtain the following solutions, via the symmetry (14):

$$\begin{aligned} y(x) &= \frac{1}{\sum_{m=0}^\infty x^n \sum_{m=-n-1}^{n+1} c_{n+1,m} x^{m\sigma}} = \frac{x^\sigma / c_{1,-1}}{1 + \frac{c_{10}}{c_{1,-1}} x^\sigma + \frac{c_{11}}{c_{1,-1}} x^{2\sigma} + \sum_{n=1}^\infty x^n \sum_{m=-n}^{n+2} \frac{c_{n+1,m-1}}{c_{1,-1}} x^{m\sigma}} \\ &= \frac{1}{c_{1,-1}} x^\sigma \sum_{n=0}^\infty x^n \sum_{m=-n}^\infty b_{nm} x^{m\sigma}, \quad b_{nm} \in \mathbf{C} \end{aligned}$$

Namely

$$y(x) = \sum_{n=1}^\infty x^n \sum_{m=-n}^\infty d_{nm} x^{m\rho}, \quad \rho = 1 - \sigma, \quad 0 < \Re \rho < 1, \quad d_{nm} \in \mathbf{C} \quad (34)$$

Note that $1 + \rho$ does not belong to the set (8). If we do the explicit computation above, we find that

$$d_{nm} = 0 \quad \text{for } m > n$$

Thus, (34) is a solution (22).

(*) We have re-obtained the class $\mathcal{A}_0$ of [6].

Note: To verify that $d_{nm} = 0$ for $m > n$, one can do the explicit computation of the d_{nm} by expanding $x/y(x)$ and computing the coefficients. Note that to get the correct terms in the truncation $\sum_{n=1}^N x^n \sum_{m=-n}^n d_{nm} x^{m\rho}$, one needs to start from (22) truncated at $\sum_{n=1}^{N+3} x^n \sum_{m=-n}^n c_{nm} x^{m\sigma}$. One can also conclude directly from the form (34) that $d_{nm} = 0$ for $m > n$. Actually, such a form, substituted into (PVI), implies $d_{nm} = 0$ for $m > n$. Namely, we always have that:

$$y(x) = \sum_{n=1}^\infty x^n \sum_{m=-\infty}^\infty d_{nm} x^{m\rho} \quad + \quad \text{substitution into (PVI)} \quad \implies \quad y(x) = \sum_{n=1}^\infty x^n \sum_{m=-n}^n d_{nm} x^{m\rho}$$

◇
1.1b) From (24) via the symmetry (14), we obtain a three-real parameters ($\nu \in \mathbf{R}$, $\phi \in \mathbf{C}$) solution (see [12]):

$$y(x) = \left[\sum_{n=0}^{\infty} x^n \sum_{m=-n-1}^{n+1} c_{nm} e^{im\phi} x^{2im\nu} \right]^{-1} = \frac{1}{A \sin(2\nu \ln x + \phi) + B + O(x)} \quad (35)$$

where

$$O(x) = \sum_{n=1}^{\infty} x^n \sum_{m=-n-1}^{n+1} c_{nm} e^{im\phi} x^{2im\nu}, \quad A = -\sqrt{\frac{\alpha}{2\nu^2} + B^2}, \quad B = \frac{2\nu^2 + \gamma - \alpha}{4\nu^2}$$

The expansion is convergent. All c_{nm} , are rational functions of ν , and $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}$. If $\arg x$ is bounded (namely, when we considered $y(x)$ as a branch), then (35) admits two infinite sequences of *movable poles* in the neighborhood of $x = 0$, asymptotically distributed along two rays and accumulating at $x = 0$ (see [13]). One can do a formal (non convergent) expansion, when x is far from the poles, as

$$y(x) = \frac{1}{A \sin(2\nu \ln x + \phi) + B} + \delta_\nu x^{2i\nu} \left\{ 1 + \sum_{n=0}^{\infty} x^n \sum_{m=-n}^{\infty} \delta_{nm} x^{2im\nu} \right\}, \quad \delta_\nu, \delta_{nm} \in \mathbf{C} \quad (36)$$

(*) The class of solutions (35) coincide with the union $(\mathcal{B}_0^+ \cup \mathcal{B}_0^-) \cup (\mathcal{B}_7^+ \cup \mathcal{B}_7^-)$ of [6], where a formal expansion, in a form equivalent to (36), is given.

◇
2.1) The symmetry (14) applied to case 2) gives one parameter ($a \in \mathbf{C}$) solutions, defined when at least one of $\boxed{\sqrt{2\alpha} + \sqrt{2\gamma} \notin \mathbf{Z}}$ or $\boxed{\sqrt{2\alpha} - \sqrt{2\gamma} \notin \mathbf{Z}}$ (namely $\theta_\infty + \theta_1 \notin \mathbf{Z}$ or $\theta_\infty - \theta_1 \notin \mathbf{Z}$). Let $\omega \in \Omega_{\alpha\gamma}$. We distinguish two cases, $\alpha \neq 0$ and $\alpha = 0$.

◇ Case $\boxed{\alpha \neq 0}$.

– **2.1a)** When $\Re \sigma > 0$ in (26) we obtain the following, via the symmetry transformation (14):

$$y(x) = \frac{1}{c_{10} \left(1 + \frac{c_{11}}{c_{10}} x^\sigma + \sum_{n=1}^{\infty} x^n \sum_{m=0}^{n+1} \frac{c_{n+1,m}}{c_{10}} x^{m\sigma} \right)} \Big|_{\sigma \mapsto \omega} = \sum_{n=0}^{\infty} x^n \sum_{m=0}^{\infty} b_{nm} x^{m\omega},$$

where b_{01} is the arbitrary constants, and the other coefficients are rational functions of $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}, \sqrt{1-2\delta}$ and b_{01} . In particular $b_{00} = \frac{\sqrt{\alpha} \pm \sqrt{\gamma}}{\sqrt{\alpha}}$. With the substitution $\omega = \rho + 1$, the above becomes

$$y(x) = \frac{\sqrt{\alpha} + (-)^k \sqrt{\gamma}}{\sqrt{\alpha}} + \sum_{n=1}^{\infty} x^n \sum_{m=0}^n d_{nm} (ax^\rho)^m = \sum_{l=0}^{\infty} y_l(x) (ax^\rho)^l \quad (37)$$

$$\rho + 1 \in \Omega_{\alpha\gamma}^k \text{ given by (7), } k = 1, 2, \Rightarrow \Re \rho > -1$$

Here a is the arbitrary constant, $d_{nm} \in \mathbf{C}$ are rational functions of $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}$, and $\sqrt{1-2\delta}$. The $y_l(x) = O(x^l)$'s are Taylor series. In particular

$$y_0(x) = \frac{\sqrt{\alpha} \pm \sqrt{\gamma}}{\sqrt{\alpha}} + \sum_{n=1}^{\infty} d_{n0} x^n, \quad \alpha \neq 0 \quad (38)$$

Note that $y_1(x)$ can be directly obtained from (27) by the symmetry (14). Its convergence is proved in [19].

(*) In terms of the classification of [6], the class of solutions (37) coincides with the union of the class $\mathcal{B}_6$ with the subclass of $\mathcal{B}_1 \cup \mathcal{B}_2$ which does not contain logarithmic coefficients.

◇

– **2.1b)** When $\Re\sigma = 0$ in (26), we obtain through (14)

$$y(x) = \left\{ \sum_{n=0}^{\infty} x^n \tau_n(x) \right\}^{-1}, \quad \tau_n(x) := \sum_{m=0}^{n+1} c_{n+1,m} (ax^{2i\nu})^m, \quad \nu \in \mathbf{R}, \quad \nu \neq 0 \quad (39)$$

Here $2i\nu = \omega \in \Omega_{\alpha\beta}$. Note that $\Omega_{\alpha\beta}$ must contain an imaginary element. We have two solutions above with $\omega = \pm(\sqrt{2\alpha} + \sqrt{2\gamma})$, when $\sqrt{2\alpha} + \sqrt{2\gamma}$ is imaginary, or we have two solutions with $\omega = \pm(\sqrt{2\alpha} - \sqrt{2\gamma})$, when $\sqrt{2\alpha} - \sqrt{2\gamma}$ is imaginary. In particular

$$\tau_0(x) = \frac{\sqrt{\alpha}}{\sqrt{\alpha} + (-)^k \sqrt{\gamma}} + ax^{2i\nu} = \frac{\sqrt{\alpha}}{\sqrt{\alpha} + (-)^k \sqrt{\gamma}} + \left\{ \cos(2\nu \ln x + \phi) + i \sin(2\nu \ln x + \phi) \right\},$$

$$\phi = \ln a, \quad 2i\nu \in \Omega_{\alpha\gamma}^k = \left\{ (\sqrt{2\alpha} + (-)^k \sqrt{2\gamma}), -(\sqrt{2\alpha} + (-)^k \sqrt{2\gamma}) \right\} \subset i\mathbf{R}, \quad k = 1, 2.$$

The coefficients $c_{n+1,m}$'s are rational functions of $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}$ and $\sqrt{1-2\delta}$. The denominator $\sum_{n=0}^{\infty} x^n \tau_n(x)$ may vanish, so $y(x)$ may have movable poles. In a neighborhood of $x = 0$ they are asymptotically distributed along the ray of the zeros of $\tau_0(x)$. The latter are given by

$$|x| = \exp \left\{ \frac{1}{2\nu} \left(\arg \frac{c_{10}}{a} + (2l+1)\pi \right) \right\}, \quad \arg x = -\frac{1}{2\nu} \ln \left| \frac{c_{10}}{a} \right|$$

$$c_{10} = \frac{\sqrt{\alpha}}{\sqrt{\alpha} \pm \sqrt{\gamma}}, \quad l \in \mathbf{Z}, \quad l < 0$$

Far from the line of the poles, we can write

$$y(x) = \frac{1}{\tau_0(x)} \left[1 - \frac{\tau_1(x)}{\tau_0(x)} + O(x^2) \right], \quad \frac{\tau_1(x)}{\tau_0(x)} = O(x) \quad (40)$$

(*) The class of solution (39), when formally expanded from (40), coincides with the union $(\mathcal{B}_1^+ \cup \mathcal{B}_1^-) \cup (\mathcal{B}_2^+ \cup \mathcal{B}_2^-) \cup (\mathcal{B}_6^+ \cup \mathcal{B}_6^-)$ of [6].

◇

– **2.1c)** When $-1 < \Re\sigma < 0$ in (26), we obtain a solution for $-1 < \Re\omega < 0$ of the form

$$y(x) = \frac{\frac{1}{c_{11}} x^{-\omega}}{1 + \frac{c_{10}}{c_{11}} x^{-\omega} + \sum_{n=1}^{\infty} x^n \sum_{m=-1}^n \frac{c_{n+1,m+1}}{c_{11}} x^{m\omega}}$$

$$= ax^{-\omega} \sum_{n=0}^{\infty} x^n \sum_{m=-n}^{\infty} b_{nm} x^{-m\omega} = \sum_{n=1}^{\infty} x^n \sum_{m=-n}^{\infty} d_{nm} x^{m(1+\omega)}$$

where $a = 1/c_{11}$ is an arbitrary constant. This should coincide with a solution (22) with $\sigma = 1 + \omega$, $0 < \Re\sigma < 1$, namely

$$y(x) = \sum_{n=1}^{\infty} x^n \sum_{m=-n}^n d_{nm} x^{m(1+\omega)} \sim ax^{-\omega}$$

(*) In terms of the classification of [6], we have obtained solutions in $\mathcal{A}_0$.

◇—————◇

◇ **Case** $\boxed{\alpha = 0}$.

◇—————◇

– **2.1d**) The symmetry (14) applied to case 2) when $\beta = 0$ yields the case $\alpha = 0$:

$$y(x) = \frac{1}{c_{11}x^\sigma + \sum_{n=1}^{\infty} x^n \sum_{m=1}^{n+1} c_{n+1,m} x^{m\sigma}} = \frac{1}{c_{11}x^\sigma \left(1 + \sum_{n=1}^{\infty} x^n \sum_{m=0}^n \frac{c_{n+1,m+1}}{c_{11}} x^{m\sigma}\right)} \Big|_{\sigma \mapsto \omega}$$

Namely

$$\boxed{y(x) = \frac{1}{a} x^{-\omega} \left(1 + \sum_{n=1}^{\infty} x^n \sum_{m=0}^n d_{nm} (ax^\omega)^m\right) = \frac{1}{a} x^{-\omega} \sum_{k=0}^{\infty} y_k(x) (ax^\omega)^k} \quad (41)$$

$$\omega = \sqrt{2\gamma} \operatorname{sgn}(\Re\sqrt{2\gamma}) \quad \text{or} \quad \omega \in \{\sqrt{2\gamma}, -\sqrt{2\gamma}\} \quad \text{if} \quad -1 < \Re\sqrt{2\gamma} < 1; \quad \omega \notin \mathbf{Z}$$

Here a is the free parameter and $\Re\omega > -1$. d_{nm} are rational functions of $\sqrt{\alpha}, \sqrt{\beta}, \sqrt{\gamma}$ and $\sqrt{1-2\delta}$. $y_l(x) = O(x^l)$ are Taylor series, and $y_0(x)$ is monic, namely $y_0(x) = 1 + O(x)$. The above $|y(x)|$ diverges for $x \rightarrow 0$ when $\Re\omega > 0$. When $\sqrt{2\gamma} \in i\mathbf{R}$, it becomes the special case $\alpha = 0$ of (39).

Note that when $-1 < \Re\omega < 0$, we obtain solutions of type (22). On the other hand, we obtain truly new behaviors (divergent, or oscillatory and non vanishing) when

$$\boxed{\omega = \sqrt{2\gamma} \operatorname{sgn}(\Re\sqrt{2\gamma}) \notin \mathbf{Z}, \quad \text{or} \quad \omega \in \{\sqrt{2\gamma}, -\sqrt{2\gamma}\} \subset i\mathbf{R} \text{ when } \gamma < 0} \quad (42)$$

(*) The class of solution (41), with restriction (42), coincides with $\mathcal{C}_0^\infty \cup \mathcal{B}_8 \cup \mathcal{B}_9$ of [6].

◇—————◇

3.1) The symmetry (14) applied to case 3), gives solutions existing when $\boxed{\sqrt{2\alpha} + \sqrt{2\gamma} = N}$ or $\boxed{\sqrt{2\alpha} - \sqrt{2\gamma} = N}$ (namely $\theta_\infty - 1 + \theta_1 = N$, or $\theta_\infty - 1 - \theta_1 = N$), $N \in \mathbf{Z}$:

$$\boxed{y(x) = \sum_{n=0}^{|N|-1} b_n x^n + \left(a + b_N \ln x\right) x^{|N|} + \sum_{n=|N|+1}^{\infty} P_n(\ln x; a) x^n, \quad N \neq 0} \quad (43)$$

$$y(x) = \frac{1}{(a \pm \sqrt{2\alpha} \ln x) + \sum_{n=1}^{\infty} P_n(\ln x; a) x^n} = \pm \frac{1}{\sqrt{2\alpha} \ln x} \left(1 \mp \frac{1}{\sqrt{2\alpha} \ln x} + O\left(\frac{1}{\ln^2 x}\right) \right), \quad \begin{matrix} N=0 \\ \alpha = \gamma \neq 0 \end{matrix}$$

if $\boxed{\{\sqrt{-2\beta} + \sqrt{1-2\delta}, \sqrt{-2\beta} - \sqrt{1-2\delta}\} \cap \{|N| - 1, 1 - |N|\} \neq \emptyset}$, namely $\{\theta_0 + \theta_x, \theta_0 - \theta_x\} \cap \{|N| - 1, 1 - |N|\} \neq \emptyset$, or if $\boxed{2\alpha = N^2}$ (namely $\theta_\infty = N + 1$). The expansions are

$$y(x) = \sum_{n=0}^{|N|-1} b_n x^n + a x^{|N|} + \sum_{n=|N|+1}^{\infty} b_n(a) x^n, \quad N \neq 0 \quad (45)$$

$$y(x) = a + (1-a)(\delta - \beta)x + \sum_{n=2}^{\infty} b_n(a)x^n, \quad N=0, \quad \alpha = \gamma = 0 \quad (46)$$

We observe that the case (45) with $N = 1$ gives two subcases:

$$y(x) = \frac{1}{\sqrt{2\alpha}} + ax + O(x^2), \quad \alpha \neq 0, \quad (\sqrt{2\alpha} \pm \sqrt{2\gamma})^2 = 1, \quad 1 - 2\delta + 2\beta = 0 \quad (47)$$

and

$$y(x) = 1 + ax + \frac{1}{4}a(1 + 4a + 2\beta - 2\delta)x^2 + O(x^3), \quad \alpha = \frac{1}{2}, \quad \gamma = 0$$

The latter is equivalent to (46) via a birational transformation.

Solutions (46), (47) and (38) are basic Taylor solutions, which generate all the other Taylor expansions by the action of the birational transformations. They are obtained in [10] making use of the matching procedure with non Fuchsian IN and OUT reductions of (1). They are also derived in [19].

(*) In terms of the classification of [6], the class of the solutions (43) is the subclass of $\mathcal{B}_1 \cup \mathcal{B}_2$ with logarithmic coefficients. The class of the solutions (44) coincides with the class $\mathcal{B}_4 \cup \mathcal{B}_5$. The sub case (46) gives the class $\mathcal{B}_{10}$ of [6].

4.1) Symmetry (14) applied to 4) gives the following solutions defined for $\alpha \neq \gamma$ (namely $\theta_\infty \pm \theta_1 \neq 1$):

$$y(x) = \frac{1}{\frac{\gamma-\alpha}{2}(a+\ln x)^2 + \frac{\alpha}{\alpha-\gamma}} + \sum_{n=1}^{\infty} P_{n+1}(\ln x)x^n = \frac{2}{(\gamma-\alpha)\ln^2 x} \left[1 - \frac{2a}{\ln x} + O\left(\frac{1}{\ln^2 x}\right) \right], \quad \alpha \neq \gamma$$

(*) In terms of the classification of [6], this is the class $\mathcal{B}_3$.

(*) **Final comparison with the results of paper [6].** We have shown that the set of the behaviors at $x = 0$ obtained in the framework of the isomondromy deformation method, in particular using the

method of [17] and the matching procedure of [10], coincides with the set of expansions at $x = 0$ obtained by power geometry in [6]. Summarizing, at $x = 0$, we have:

Behaviors of [17],[22],[9],[10],[11],[12]		39 families of [6]
1) behavior (22), with $\Re\sigma \neq 0$ [This includes cases 1.1a), 2) with $-1 < \Re\sigma < 1$, and 2.1c)]	=	$\mathcal{A}_0$
1.1b) behavior (35), and 1) behavior (24)	=	$\mathcal{B}_0^+ \cup \mathcal{B}_0^- \cup \mathcal{B}_7^+ \cup \mathcal{B}_7^-$ and image ¹
2.1a) behavior (37), and 2) behavior (26) if $\Re\sigma > 0$	=	$\mathcal{B}_1(\text{non log}) \cup \mathcal{B}_2(\text{non log}) \cup \mathcal{B}_6$ and image
2.1b) behavior (39), and 2) behavior (26) if $\Re\sigma = 0$	=	$\mathcal{B}_1^+ \cup \mathcal{B}_1^- \cup \mathcal{B}_2^+ \cup \mathcal{B}_2^- \cup \mathcal{B}_6^+ \cup \mathcal{B}_6^-$ and image
2.1d) behavior (41), and 2) behavior (26) if $\beta = 0$	=	$\mathcal{C}_0^\infty \cup \mathcal{B}_8 \cup \mathcal{B}_9$ and image
3.1), and 3)	=	$\mathcal{B}_1(\text{log}) \cup \mathcal{B}_2(\text{log}) \cup \mathcal{B}_4 \cup \mathcal{B}_5 \cup \mathcal{B}_{10}$ and image
4.1), and 4)	=	$\mathcal{B}_3$ and image

¹“image” means the image of the families indicated through (14).

In [6] a classification of actual, asymptotic or formal expansions which satisfy an ODE is given. In terms of it, we have that:

- Solutions (22) and (26) for $\Re\sigma \neq 0$, (37) for $\Re\rho \neq 0$, (41) for $\Re\omega \neq 0$, and the Taylor series (30), (31), (45), (46), (27) and (38) have *power expansions*.
- Solutions (28), (29), (33) and (43) have *power logarithmic expansions*.
- Solutions (44) and (48), when expanded in powers of $(\ln x)^{-1}$, have *complicated expansions*.
- Solutions (24) (namely, (22) for $\Re\sigma = 0$) and (26) for $\Re\sigma = 0$, (37) for $\Re\rho = 0$, and (41) for $\Re\omega = 0$ have *semi exotic expansions*.
- Solutions (35) and (39), when formally expanded (like in (36)), have *exotic expansions*.

8 Details on the Parameterization in terms of monodromy data. Connection Problem

As already explained, to every branch $y(x)$ of a solution of PVI, a system (1) is associated, and thus to $y(x)$ the monodromy group of the system is associated, namely, monodromy data are associated (definition of monodromy data is in section 4).

It may well happen that to a family of solutions $y(x)$ the same monodromy group is associated. An example, which is found in [11], is a one parameter $a \in \mathbf{C}$ class of solutions existing when

$$\alpha = \gamma = 0, \quad \delta = \frac{1}{2}, \quad \beta = -2p^2; \quad p \in \mathbf{Z}, \quad p \neq 0$$

Their critical behavior is power logarithmic or inverse power logarithmic at the three critical points:

$$y(x) \sim \begin{cases} x [1 - p^2 (\ln x + a)^2], & x \rightarrow 0, \\ 1 - p^2 \left(\ln \frac{1}{x} + \rho_\infty \right)^2, & x \rightarrow \infty, \\ 1 - \frac{1}{p^2 (\ln(1-x) + \rho_1)^2}, & x \rightarrow 1. \end{cases} \quad (49)$$

where:

$$\rho_\infty = \frac{\pi(4 \ln 2 - 1 + a)}{\pi - i(4 \ln 2 - 1 + a)} - 2 \ln 2 + 1, \quad \rho_1 = \frac{\pi^2}{4 \ln 2 - 1 + a} - \ln 2 + 1.$$

The monodromy group associated to (49) is independent of the parameter a , and is generated by:

$$M_0 = I, \quad M_x = \begin{pmatrix} 1 & 2\pi i \\ 0 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 1 - \frac{8i}{\pi} \ln 2 & -\frac{32i}{\pi} (\ln 2)^2 \\ \frac{2i}{\pi} & 1 + \frac{8i}{\pi} \ln 2 \end{pmatrix}.$$

In particular, $\text{Tr}(M_0 M_x) = \text{Tr}(M_0 M_1) = 2$ and $\text{Tr}(M_1 M_x) = -2$.

Cases like the above example occur for special values of the coefficients of PVI. In general to $y(x)$ is associated a point in the space of monodromy data and to this point no other branch than $y(x)$ corresponds. In section 4 we have seen that this occurs if and only if none of M_0 , M_x , M_1 and M_∞ is the identity. Moreover the values p_0 , p_x , p_1 , p_∞ , p_{0x} , p_{x1} and p_{01} are good coordinates of the subspace of monodromy data such that the group $\langle M_0, M_x, M_1 \rangle$ is not reducible and none of the matrices M_0 , M_1 , M_x and M_∞ is the identity.

Let the branch $y(x)$ depend on (at most) two integration constants c_1 and c_2 :

$$y(x) = y(x, c_1, c_2)$$

for example, in case of (22), $(c_1, c_2) = (\sigma, c_{11})$ (or (σ, a) in the table). When none of M_0 , M_x , M_1 and M_∞ is the identity, and $\langle M_0, M_x, M_1 \rangle$ is not reducible, $y(x)$ is uniquely associated to a point in the space of monodromy data, and the integration constants c_1, c_2 are uniquely determined functions:

$$c_i = c_i(p_0, p_x, p_1, p_\infty, p_{0x}, p_{x1}, p_{01}), \quad i = 1, 2 \quad (50)$$

[Note: $(p_0, p_x, p_1, p_\infty)$, or equivalently $(\theta_0, \theta_x, \theta_1, \theta_\infty)$, are given by the coefficients of PVI. Only two of p_{0x}, p_{x1}, p_{01} are independent, because they are linked by the relation $M_1 M_x M_0 = M_\infty$, which is a cubic in terms of the p_μ and p_{ij} . Thus, two are the genuine free complex parameters].

In case c_1, c_2 appear as the constants in the explicit formula of the critical behavior of $y(x)$ at one critical point, then the method of monodromy preserving deformations allows to *compute (50) explicitly*. For a given point in the space of monodromy data, namely for a given $y(x)$, the parameterization can be done at each of the critical points $x = 0, 1, \infty$. In this way *the connection problem is solved*.

In order to compute the monodromy data of (1) associated to a critical behavior of $y(x)$ at $x = 0$, we proceed as follows. *Once the matching $\Psi_{OUT} \leftrightarrow \Psi_{IN}$ in (18) has been completed* – and thus the critical behavior of $y(x)$ has been determined – we have to match Ψ_{OUT} with a fundamental solution Ψ of (1) at $\lambda = \infty$, and we have to match Ψ_{IN} with *the same* Ψ in another region of the λ -plane, typically around $\lambda = 0$ or x . If this is done, then M_1 of Ψ coincides with M_1^{OUT} of Ψ_{OUT} , while M_0 and M_x coincide with M_0^{IN} and M_x^{IN} of Ψ_{IN} .

The crucial point is that we are able to compute the monodromy matrices M_1^{OUT} , M_0^{IN} and M_x^{IN} of the reduced OUT and IN-systems exactly, namely that we are able to solve these systems in terms of linear special functions.

The entries of the monodromy matrices so computed, depend on the integration constants c_1, c_2 , namely they depend on the free parameters which govern the critical behavior of $y(x)$. In case $y(x)$ is uniquely associated to the monodromy matrices just computed (namely, none of M_0 , M_x , M_1 and M_∞ is the identity, and $\langle M_0, M_x, M_1 \rangle$ is not reducible), we calculate the traces of $M_i M_j$

$$\begin{cases} p_{0x} = p_{0x}(c_1, c_2, \theta_0, \theta_x, \theta_1, \theta_\infty) \\ p_{x1} = p_{x1}(c_1, c_2, \theta_0, \theta_x, \theta_1, \theta_\infty) \\ p_{01} = p_{01}(c_1, c_2, \theta_0, \theta_x, \theta_1, \theta_\infty) \end{cases} \quad (51)$$

These are classical functions of $(c_1, c_2, \theta_0, \theta_x, \theta_1, \theta_\infty)$. This follows from the fact that they are obtained by means of asymptotic reduction of (1) to the systems (17) and (16), which can be solved in terms of

classical functions (for definition of classical functions see [23], [24] and [25], or the introduction of [12]) The system (51) can be solved in c_1 and c_2 , and we obtain (50). Moreover, the functions in (50) are classical of their arguments. They are the parametrization formulas of section 3.

It is enough to give the parameterization at $x = 0$, because the critical behaviors and the parameterizations at $x = 1$ and ∞ can be obtained via (12) and (13) provided we change the parameters in the parameterization formulas according to

$$\begin{cases} p'_{01} &= -p_{01} - p_{0x}p_{x1} + p_{\infty}p_x + p_1p_0 \\ p'_{0x} &= p_{x1} \\ p'_{x1} &= p_{0x} \end{cases} \quad \text{or} \quad \begin{cases} p_{01} &= -p'_{01} - p'_{0x}p'_{x1} + p'_{\infty}p'_x + p'_1p'_0 \\ p_{0x} &= p'_{1x} \\ p_{x1} &= p'_{0x} \end{cases} \quad (52)$$

for (12), and

$$\begin{cases} p'_{0x} &= -p_{01} - p_{0x}p_{x1} + p_{\infty}p_x + p_0p_1 \\ p'_{01} &= p_{0x} \\ p'_{1x} &= p_{1x} \end{cases} \quad \text{or} \quad \begin{cases} p_{01} &= -p'_{0x} - p'_{01}p'_{x1} + p'_{\infty}p'_1 + p'_0p'_x \\ p_{0x} &= p'_{01} \\ p_{1x} &= p'_{1x} \end{cases} \quad (53)$$

for (13).

Note also that it is enough to compute the parametrization formulas for the cases $\mathbf{N}$), $\mathbf{N}=1,2,3,4$, of section 7. One can obtain the parameterization for the solutions in cases $\mathbf{N.1}$), from that of the equivalent cases $\mathbf{N}$) [equivalent via symmetry (14)] . As it is explained in [12], section 4, one needs to make the substitution

$$(p_0, p_x, p_1, p_{\infty}; p_{0x}, p_{01}, p_{x1}) \mapsto (-p_{\infty}, p_1, p_x, -p_0; -p_{0x}, -p_{01}, p_{x1}), \quad \sigma \mapsto 1 - \sigma \quad (54)$$

in the formulas of cases $\mathbf{N}$).

9 Further Details on the Matching to obtain Monodromy Data

The standard choice of a fundamental solution of (1) is

$$\Psi(\lambda) = \begin{cases} [I + O(\frac{1}{\lambda})] \lambda^{-\frac{\theta_{\infty}}{2}\sigma_3} \lambda^{R_{\infty}}, & \lambda \rightarrow \infty; \\ \psi_0(x)[I + O(\lambda)] \lambda^{\frac{\theta_0}{2}\sigma_3} \lambda^{R_0} C_0, & \lambda \rightarrow 0; \\ \psi_x(x)[I + O(\lambda - x)] (\lambda - x)^{\frac{\theta_x}{2}\sigma_3} (\lambda - x)^{R_x} C_x, & \lambda \rightarrow x; \\ \psi_1(x)[I + O(\lambda - 1)] (\lambda - 1)^{\frac{\theta_1}{2}\sigma_3} (\lambda - 1)^{R_1} C_1, & \lambda \rightarrow 1; \end{cases} \quad (55)$$

Here $\psi_0(x)$, $\psi_x(x)$, $\psi_1(x)$ are the diagonalizing matrices of $A_0(x)$, $A_1(x)$, $A_x(x)$ respectively. They are defined by multiplication to the right by arbitrary diagonal matrices, possibly depending on x . C_{ν} , $\nu = \infty, 0, x, 1$, are invertible *connection matrices*, independent of x [18]. Each R_{ν} , $\nu = \infty, 0, x, 1$, is also independent of x , and:

$$R_{\nu} = 0 \text{ if } \theta_{\nu} \notin \mathbf{Z}, \quad R_{\nu} = \begin{cases} \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}, & \text{if } \theta_{\nu} > 0 \text{ integer} \\ \begin{pmatrix} 0 & 0 \\ * & 0 \end{pmatrix}, & \text{if } \theta_{\nu} < 0 \text{ integer} \end{cases}$$

If $\theta_i = 0$, $i = 0, x, 1$, then R_i is to be considered the Jordan form $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ of A_i .

(*) **Matching $\Psi \leftrightarrow \Psi_{OUT}$:**

$\lambda = \infty$ is a fuchsian singularity of the OUT-system (16), with residue $-A_\infty/\lambda$. Therefore, we can always find a fundamental matrix solution with behavior:

$$\Psi_{OUT}^{Match} = \left[I + O\left(\frac{1}{\lambda}\right) \right] \lambda^{-\frac{\theta_\infty}{2}\sigma_3} \lambda^{R_\infty}, \quad \lambda \rightarrow \infty.$$

This solution matches with Ψ . Also $\lambda = 1$ is a fuchsian singularity of (16). Therefore, we have:

$$\Psi_{OUT}^{Match} = \psi_1^{OUT}(x) [I + O(\lambda - 1)] (\lambda - 1)^{\frac{\theta_1}{2}\sigma_3} (\lambda - 1)^{R_1} C_1^{OUT}, \quad \lambda \rightarrow 1;$$

Here C_1^{OUT} is a suitable connection matrix. $\psi_1^{OUT}(x)$ is the matrix that diagonalizes the leading terms of $A_1(x)$. Therefore, $\psi_1(x) \sim \psi_1^{OUT}(x)$ for $x \rightarrow 0$. As a consequence of isomonodromicity, R_1 is the same of Ψ .

As a consequence of the matching $\Psi \leftrightarrow \Psi_{OUT}^{Match}$, the monodromy of Ψ at $\lambda = 1$ is:

$$M_1 \equiv M_1^{OUT} = C_1^{-1} \exp\{i\pi\theta_1\sigma_3\} \exp\{2\pi i R_1\} C_1, \quad \text{with } C_1 \equiv C_1^{OUT}.$$

We finally need an invertible connection matrix C_{OUT} to connect Ψ_{OUT}^{Match} with the solution Ψ_{OUT} which was matching with Ψ_{IN} . Namely

$$\Psi_{OUT}^{Match} = \Psi_{OUT} C_{OUT}.$$

(*) **Matching $\Psi \leftrightarrow \Psi_{IN}$:**

As a consequence of the matching $\Psi \leftrightarrow \Psi_{OUT}^{Match}$, we have to choose the IN-solution of (17) which matches with Ψ_{OUT}^{Match} . This is

$$\Psi_{IN}^{Match} := \Psi_{IN} C_{OUT}.$$

Now, $\lambda = 0, x$ are fuchsian singularities of the IN-system. Therefore:

$$\Psi_{IN}^{Match} = \begin{cases} \psi_0^{IN}(x) [I + O(\lambda)] \lambda^{\frac{\theta_0}{2}\sigma_3} \lambda^{R_0} C_0^{IN}, & \lambda \rightarrow 0; \\ \psi_x^{IN}(x) [I + O(\lambda - x)] (\lambda - x)^{\frac{\theta_x}{2}\sigma_3} (\lambda - x)^{R_x} C_x^{IN}, & \lambda \rightarrow x; \end{cases}$$

The above hold for fixed small $x \neq 0$. Here C_0^{IN} and C_x^{IN} are suitable connection matrices. $\psi_0^{IN}(x)$ and $\psi_x(x)^{IN}$ are diagonalizing matrices of the leading terms of $A_0(x)$ and $A_x(x)$. For $x \rightarrow 0$ they match with $\psi_0(x)$ and $\psi_x(x)$ of Ψ in (55). On the other hand, as a consequence of isomonodromicity, the matrices R_0 and R_x are the same of Ψ . The above Ψ_{IN}^{Match} has the same behavior of Ψ at $\lambda \rightarrow 0$ and $\lambda \rightarrow x$; moreover, it is an approximation of Ψ for x small. The matrices C_0^{IN} , C_x^{IN} are independent of x . So, the matching $\Psi \leftrightarrow \Psi_{IN}$ is realized and the connection matrices C_0 and C_x coincide with C_0^{IN} , C_x^{IN} respectively. As a result, we obtain the monodromy matrices for Ψ :

$$\begin{aligned} M_0 &\equiv M_0^{IN} = C_0^{-1} \exp\{i\pi\theta_0\sigma_3\} \exp\{2\pi i R_0\} C_0, & C_0 &\equiv C_0^{IN}, \\ M_x &\equiv M_x^{IN} = C_x^{-1} \exp\{i\pi\theta_x\sigma_3\} \exp\{2\pi i R_x\} C_x, & C_x &\equiv C_x^{IN}. \end{aligned}$$

The matrices C_1^{OUT} , C_{OUT} , C_0^{IN} and C_x^{IN} can be computed because we are able to solve exactly (i.e. in terms of special functions) the reduced systems OUT and IN. Details are technically complicated, and the reader is referred to [10], [11].

10 Conclusions

We have given the table of critical behaviors at $x = 0$ of PVI-transcendents, obtained with the method of monodromy preserving deformations (essentially, the Jimbo's procedure [17] and the matching procedure [10]). From the table, all the behaviors at $x = 1$ and ∞ can be computed via a birational transformation (12) and (13). We have explained in details the meaning of the behaviors, and reviewed how they were obtained. We have also given the parameterization in terms of monodromy data of the basic critical behaviors. The parameterization allows to solve the connection problem. We have explained that the table of critical behaviors is essentially complete. We have shown that the expansions obtained by power geometry in [6] coincide with the behaviors obtained by the monodromy preserving deformation method.

This paper is aimed at providing a synthetic (and hopefully useful) tabulation of both critical behaviors and parametrization formulas, which so far has been missing.

Acknowledgments: I thank the Korea Institute of Advanced Study (KIAS, Seoul 130-722, Republic of Korea) for providing computing resources (Abacus System) for this work.

References

- [1] A.V. Andreev, A.V. Kitaev: *Connection formulas for asymptotics of the fifth Painlevé transcendent on the real axis*. Nonlinearity, **13**, (2000), 1801-1840.
- [2] Boalch, P. From Klein to Painlevé, via Fourier, Laplace and Jimbo. Proc. London Math. Soc. (2005) 90(1): 167-208.
- [3] Y.V. Brezhnev: *τ -function solution of the Sixth Painlevé Transcendent*. Theoretical and Mathematical Physics, **161**(3), (2009), 1616 - 1633.
- [4] A.D. Bruno: *Power Geometry in Algebraic and Differential Equations*. Nauka, Moscow, 1979; English transl., North-Holland, Amsterdam, 2000. MR1773512 (2002c:37071)
- [5] A.D. Bruno: *Asymptotic behaviour and expansion of a solution of an ordinary differential equation*. Russian Math. Surveys **59**, (2004), 429-480.
- [6] A.D. Bruno, I.V. Goryuchkina: *Asymptotic Expansions of Solutions of the Sixth Painlevé Equation*. Moscow Math. Soc., **S0077-1554(2010)00186-0**, (2010), 1-104.
- [7] A. Fokas, A. Its, A. Kapaev, V. Novokshenov: *Painlevé Transcendents: The Riemann-Hilbert Approach*. AMS (2006).
- [8] D. Guzzetti *On the Critical Behavior, the Connection Problem and the Elliptic Representation of a Painlevé 6 Equation*. Mathematical Physics, Analysis and Geometry, **4**, (2001), 293-377.
- [9] D. Guzzetti: *The Elliptic Representation of the General Painlevé VI Equation*, Comm. Pure Appl. Math. LV, (2002), 1280-1363.
- [10] D. Guzzetti: *Matching procedure for the sixth Painlevé equation*, J. Phys. A: Math. Gen. **39** (2006), 11973-1231
- [11] D. Guzzetti: *The Logarithmic asymptotics of the sixth Painlevé equation*, J. Phys. A: Math. Theor. **41** (2008), 205201(46p)
- [12] D. Guzzetti: *Solving the sixth Painlevé equation: towards the classification of all the critical behaviors and the connection formulas*. International Mathematics Research Notices 2011 rnr071, 62 pages. doi:10.1093/imrn/rnr071
- [13] D. Guzzetti *Poles Distribution of PVI transcendents close to a critical point*, arXiv1104.5066[math.CA] (2011)

- [14] N.Hitchin: *Hypercomplex Manifolds and the Space of Framings*, The Geometric Universe, Oxford Univ. Press, (1998), 9-30.
- [15] A.R.Its, V.Y.Novokshenov: *The isomonodromy deformation method in the theory of Painlevé equations*, Springer Lecture Notes in Mathematics 1191, (1986).
- [16] K. Iwasaki: *An area-preserving action of the modular group on cubic surfaces abd the Painlevé VI equation*, Comm.Math.Phys., **242**, (2003) 185-219.
- [17] M.Jimbo: *Monodromy Problem and the Boundary Condition for Some Painlevé Trascendents*. Publ. RIMS, Kyoto Univ.,**18** (1982), 1137-1161.
- [18] M.Jimbo, T.Miwa, K.Ueno: *Monodromy Preserving Deformations of Linear Ordinary Differential Equations with Rational Coefficients (I)*. Physica **D 2** , (1981), 306-352
- [19] K.Kaneko: *Painlevé VI transcendents which are meromorphic at a fixed singularity*. Proceedings of the Japan Academy, Vol 82, Ser. A, No. 5 (2006), 71-76 .
- [20] M.Mazzocco: *Rational Solutions of the Painlevé VI Equation*, J. Phys. A: Math. Gen. **34**, (2001), 2281–2294.
- [21] K.Okamoto: *Studies on the Painlevé Equation. I – Sixth Painlevé Equation P_{VI}* , Ann. Mat. Pura Appl. **148**, (1987), 337-381.
- [22] S.Shimomura: *A Family of Solutions of a Nonlinear Ordinary Differntial Equation and its Application to Painlevé Equations (III), (V), (VI)*. J.Math. Soc. Japan, **39**, (1987), 649-662.
- [23] H.Umemura: *Painlevé Birational automorphism groups and differential equations*, Nagoya Math. J.,**119**, (1990), 1-80.
- [24] H.Umemura: *On the Irreducibility of the First Differential Equation of Painlevé*, Algebraic Geometry and Commutative Algebra in honor of Masayoshi NAGATA, Tokyo: Kinokuniya 1987, 771-789.
- [25] H.Umemura: *Second proof of the irreducibility of the First Differential Equation of Painlevé*, Nagoya Math. J.,**117**, (1990), 125-171